

Quantum deformations of the restriction of $GL_{mn}(\mathbb{C})$ -modules to $GL_m(\mathbb{C}) \times GL_n(\mathbb{C})$

Dedicated to Sri Ramakrishna

Bharat Adsul, Milind Sohoni *
Department of Computer Science and Engg., I.I.T. Bombay

K. V. Subrahmanyam †
Chennai Mathematical Institute

May 1, 2009

Abstract

In this paper, we consider the restriction of finite dimensional $GL_{mn}(\mathbb{C})$ -modules to the subgroup $GL_m(\mathbb{C}) \times GL_n(\mathbb{C})$. In particular, for a Weyl module $V_\lambda(\mathbb{C}^{mn})$ of $U_q(gl_{mn})$ we construct a representation W_λ of $U_q(gl_m) \otimes U_q(gl_n)$ such that at $q = 1$, the restriction of $V_\lambda(\mathbb{C}^{mn})$ to $U_1(gl_m) \otimes U_1(gl_n)$ matches its action on W_λ at $q = 1$. Thus W_λ is a q -deformation of the module V_λ . This is achieved by first constructing a $U_q(gl_m) \otimes U_q(gl_n)$ -module $\wedge^k$, a q -deformation of the simple $GL_{mn}(\mathbb{C})$ -module $\wedge^k(\mathbb{C}^{mn})$. We also construct the bi-crystal basis for $\wedge^k$ and show that it consists of signed subsets. Next, we develop $U_q(gl_m) \otimes U_q(gl_n)$ -equivariant maps $\psi_{a,b} : \wedge^{a+1} \otimes \wedge^{b-1} \rightarrow \wedge^a \otimes \wedge^b$. This is used as the building block to construct the general W_λ .

1 Introduction

$GL_N(\mathbb{C})$ will denote the general linear group of invertible $N \times N$ complex matrices, and $gl_N(\mathbb{C})$ its Lie algebra. Consider the group $GL_m(\mathbb{C}) \times GL_n(\mathbb{C})$ acting on X , the space of $m \times n$ -matrices with complex entries, as follows:

$$(a, b) \cdot x \rightarrow a \cdot x \cdot b^T$$

where $a \in GL_m(\mathbb{C})$, $b \in GL_n(\mathbb{C})$ and $x \in X$. Via this action, we have a homomorphism

$$\phi : GL_m(\mathbb{C}) \times GL_n(\mathbb{C}) \rightarrow GL_{mn}(\mathbb{C})$$

For a Weyl module $V_\lambda(X)$, via ϕ , we have:

$$V_\lambda(X) = \oplus_{\alpha, \beta} n_{\alpha, \beta}^\lambda V_\alpha(\mathbb{C}^m) \otimes V_\beta(\mathbb{C}^n)$$

The numbers $n_{\alpha, \beta}^\lambda$ and its properties are of abiding interest. Even the simplest question of when is $n_{\alpha, \beta}^\lambda > 0$ remains unanswered.

Our own motivation comes from the outstanding problem of P vs. NP, and other computational complexity questions in theoretical computer science (see [16]). More specifically, we look at the geometric-invariant-theoretic approach to the problem, as proposed in [13, 14]. In this approach, the

*adsul, sohoni@cse.iitb.ac.in

†kv@cmi.ac.in

general subgroup restriction problem, i.e., analysing an irreducible representation of a group G when restricted to a subgroup $H \subseteq G$, is an important step. An approach to the problem was presented in [15], via the dual notion of FRT-algebras (see, e.g., [11]); more on this later.

A useful tool in the analysis of representations of the linear groups $GL_N(\mathbb{C})$ (henceforth, just GL_N), has been the *quantizations* $U_q(gl_N)$ of the enveloping algebra of the lie algebra $gl_N(\mathbb{C})$, see [1, 3, 4, 6, 9, 10, 12]. The representation theory of $U_q(gl_N)$ mimics that of GL_N and has contributed significantly to the understanding of the diagonal embedding $GL_N \rightarrow GL_N \times GL_N$, i.e., in the tensor product of Weyl modules. This is achieved by the Hopf $\Delta : U_q(gl_N) \rightarrow U_q(gl_N) \otimes U_q(gl_N)$, a q -deformation of the diagonal embedding. However, there seems to be no quantization of $\phi : GL_m \times GL_n \rightarrow GL_{mn}$, i.e., an algebra map (also $\phi : U_q(gl_m) \otimes U_q(gl_n) \rightarrow U_q(gl_{mn})$); perhaps none exists [7].

On the other hand, we may separately construct embeddings $U_q(gl_m) \rightarrow U_q(gl_{mn})$ and $U_q(gl_n) \rightarrow U_q(gl_{mn})$ which correspond to ϕ at $q = 1$. However, the images $(U_q(gl_m))$ and $(U_q(gl_n))$ do not commute within $U_q(gl_{mn})$. This prevents the standard $U_q(gl_{mn})$ -module $V_\lambda(\mathbb{C}^{mn})$ from becoming a $U_q(gl_m) \otimes U_q(gl_n)$ -module.

This paper constructs a $U_q(gl_m) \otimes U_q(gl_n)$ -module W_λ with the following properties.

- W_λ has a weight structure which matches that of $V_\lambda(\mathbb{C}^{mn})$. Further, there is a weight-preserving bijection $W_\lambda \rightarrow V_\lambda(\mathbb{C}^{mn})$.
- The action of $U_q(gl_m) \otimes U_q(gl_n)$ on W_λ at $q = 1$ matches the action of $U_1(gl_m) \otimes U_1(gl_n)$ via the embedding $\phi : U_1(gl_m) \times U_1(gl_n) \rightarrow U_1(gl_{mn})$ on $V_\lambda(\mathbb{C}^{mn})$.

This construction is done in three steps. We first construct $U_q(gl_m) \otimes U_q(gl_n)$ -modules W_λ when $V_\lambda = \wedge^k(\mathbb{C}^{mn})$, i.e., λ is a single column shape. Next, we construct $U_q(gl_m) \otimes U_q(gl_n)$ -equivariant maps

$$\psi_{a,b} : \wedge^{a+1}(\mathbb{C}^{mn}) \otimes \wedge^{b-1}(\mathbb{C}^{mn}) \rightarrow \wedge^a(\mathbb{C}^{mn}) \otimes \wedge^b(\mathbb{C}^{mn})$$

whose co-kernel is W_λ when λ has two columns. Finally, the above map gives us straightening relations which yield the construction of general W_λ . Both, the construction of $\wedge^k(\mathbb{C}^{mn})$ and the map $\psi_{a,b}$ are deformations of the usual $U_1(gl_{mn})$ -structures, at $q = 1$.

We use the standard model for $U_q(gl_n)$ and its modules consisting of semi-standard young tableau, see, e.g., [8]. Thus a basis for $V_\lambda(\mathbb{C}^{mn})$ is identified with $SS(\lambda, mn)$, i.e., semi-standard tableau of shape λ with entries in $[mn]$.

In Section 2, we set up notation and then construct the $U_q(gl_m) \otimes U_q(gl_n)$ -modules $\wedge^k$. In the next section, we construct the abstract module W_λ for general λ . Section 4 proves some elementary properties of $U_q(gl_m) \otimes U_q(gl_n)$ -modules in the chosen basis parametrized by column tableaux. This is used for an explicit construction of $\psi_{a,b}$. In Section 5, we revert back to $\wedge^k(\mathbb{C}^{mn})$ and prove that signed column-tableaus do indeed form a bi-crystal basis for the $U_q(gl_m) \otimes U_q(gl_n)$ -action thus validating the construction in [2].

The construction in this paper has many similarities with that in [15]. Indeed, our construction of the basic subspaces $\wedge^2(\mathbb{C}^{mn})$ and $Sym^2(\mathbb{C}^{mn})$ of $\mathbb{C}^{mn} \otimes \mathbb{C}^{mn}$ is identical to that in [15]. There, these subspaces are used to construct the R -matrix and the dual algebra $GL_q(\overline{\mathbb{C}^{mn}})$ and maps $GL_q(\overline{\mathbb{C}^{mn}}) \rightarrow GL_q(\mathbb{C}^m) \otimes GL_q(\mathbb{C}^n)$. The representation theory of $GL_q(\overline{\mathbb{C}^{mn}})$ does not quite match that of the standard $GL_q(\mathbb{C}^{mn})$ and thus the construction of $V_\lambda(\mathbb{C}^{mn})$ must follow a different route. Our construction starts with the same R -matrix but bypasses the construction of $GL_q(\overline{\mathbb{C}^{mn}})$ to arrive directly at a $GL_q(\mathbb{C}^m) \otimes GL_q(\mathbb{C}^n)$ -structure for $\wedge^k(\mathbb{C}^m)$. As in [15], we have the ‘compactness’ observation, see Proposition 33. However, many other structures of [15] are as yet missing.

2 The $U_q(gl_m) \otimes U_q(gl_n)$ structure for $\wedge^k(\mathbb{C}^{mn})$

To begin, we lift *almost verbatim*, the initial parts of Section 2 of [8]. $U_q(gl_N)$ is the associative algebra over $\mathbb{C}(q)$ generated by the $4N - 2$ symbols $e_i, f_i, i = 1, \dots, N - 1$ and $q^{\epsilon_i}, q^{-\epsilon_i}, i = 1, \dots, N$ subject

to the relations:

$$\begin{aligned}
q^{\epsilon_i} q^{-\epsilon_i} &= q^{-\epsilon_i} q^{\epsilon_i} = 1, \quad [q^{\epsilon_i}, q^{\epsilon_j}] = 0 \\
q^{\epsilon_i} e_j q^{-\epsilon_i} &= \begin{cases} q e_j & \text{for } i = j \\ q^{-1} e_j & \text{for } i = j + 1 \\ e_j & \text{otherwise} \end{cases} \\
q^{\epsilon_i} f_j q^{-\epsilon_i} &= \begin{cases} q^{-1} f_j & \text{for } i = j \\ q f_j & \text{for } i = j + 1 \\ f_j & \text{otherwise} \end{cases} \\
[e_i, f_j] &= \delta_{ij} \frac{q^{\epsilon_i} q^{-\epsilon_{i+1}} - q^{-\epsilon_i} q^{\epsilon_{i+1}}}{q - q^{-1}} \\
[e_i, e_j] &= [f_i, f_j] = 0 \text{ for } |i - j| > 1 \\
e_j e_i^2 - (q + q^{-1}) e_i e_j e_i + e_i^2 e_j &= f_j f_i^2 - (q + q^{-1}) f_i f_j f_i + f_i^2 f_j = 0 \text{ when } |i - j| = 1
\end{aligned}$$

The subalgebra generated by e_i, f_i and

$$q^{h_i} = q^{\epsilon_i} q^{-\epsilon_{i+1}} \quad q^{-h_i} = q^{-\epsilon_i} q^{\epsilon_{i+1}} \quad \text{for } i = 1, \dots, N-1$$

is denoted by $U_q(sl_N)$.

The $U_q(gl_N)$ module $V_{(1^k)}$ (henceforth $\wedge^k(\mathbb{C}^N)$) is an $\binom{N}{k}$ -dimensional $C(q)$ -vector space with basis $\{v_c\}$ indexed by the subsets c of $[N]$ with k elements, i.e., by Young Tableau of shape (1^k) with entries in $[N]$. The action of $U_q(gl_N)$ on this basis is given by

$$\begin{aligned}
q^{\epsilon_i} v_c &= \begin{cases} v_c & \text{if } i \notin c \\ q v_c & \text{otherwise} \end{cases} \\
e_i v_c &= \begin{cases} 0 & \text{if } i+1 \notin c \text{ or } i \in c \\ v_d & \text{otherwise, where } d = c - \{i+1\} + \{i\} \end{cases} \\
f_i v_c &= \begin{cases} 0 & \text{if } i+1 \in c \text{ or } i \notin c \\ v_d & \text{otherwise, where } d = c - \{i\} + \{i+1\} \end{cases}
\end{aligned}$$

In order to construct more interesting modules, we use the tensor product operation. Given two $U_q(gl_N)$ -modules M, L , we can define a $U_q(gl_N)$ -structure on $M \otimes L$ by putting

$$\begin{aligned}
q^{\epsilon_i}(u \otimes v) &= q^{\epsilon_i} u \otimes q^{\epsilon_i} v \\
e_i(u \otimes v) &= e_i u \otimes v + q^{-h_i} u \otimes e_i v \\
f_i(u \otimes v) &= f_i u \otimes q^{h_i} v + u \otimes f_i v
\end{aligned}$$

Indeed, the Hopf map $\Delta : U_q(gl_N) \rightarrow U_q(gl_N) \otimes U_q(gl_N)$:

$$\Delta q^{\epsilon_i} = q^{\epsilon_i} \otimes q^{\epsilon_i}, \Delta e_i = e_i \otimes 1 + q^{-h_i} \otimes e_i, \Delta f_i = f_i \otimes q^{h_i} + 1 \otimes f_i$$

is an algebra homomorphism and makes $U_q(gl_N)$ into a bialgebra.

2.1 Some basic lemmas

We consider the $U_q(gl_{mn})$ -module $\wedge^p(\mathbb{C}^{mn})$, i.e., the homomorphism $U_q(gl_{mn}) \rightarrow \text{End}_{\mathbb{C}(q)}(\wedge^p(\mathbb{C}^{mn}))$. We gather together some lemmas on this particular action.

Lemma 1 *On the module $\wedge^p(\mathbb{C}^{mn})$, we have:*

- $e_i^2 = 0$ for all i .

- $e_i e_j e_i = 0$ whenever $|j - i| = 1$.
- $e_i f_{i+1} = e_{i+1} f_i = 0$ for all i .

We have this important combinatorial lemma:

Lemma 2 Let $\sigma = [\sigma_1, \dots, \sigma_n]$ integers such that the set $\{\sigma_1, \dots, \sigma_n\} = \{1, \dots, n\}$. Then, on the module $\wedge^p(\mathbb{C}^{mn})$, for the monomial $e_\sigma = e_{\sigma_1} \dots e_{\sigma_n}$ there exists positive integers $k_1, \dots, k_s$ such that $\sum_i k_i = n$ and

$$e_\sigma = e_{n-k_s+1} e_{n-k_s+2} \dots e_n e_{n-k_s-k_{s-1}+1} e_{n-k_s-k_{s-1}+2} \dots e_{n-k_s} \dots e_1 e_2 \dots e_{k_1}$$

An important property of the re-ordering is that either (i) the position of e_i is to the left of position of e_{i-1} or (ii) is **immediately to the right**.

Example 3 We may verify that:

$$e_2 e_6 e_7 e_3 e_5 e_1 e_4 = e_6 e_7 e_5 e_2 e_3 e_4 e_1$$

with $k_1 = 1, k_2 = 3, k_3 = 1, k_4 = 2$.

Corollary 4 Let σ be a permutation on the set $\{i, \dots, j\}$ then for the action on $\wedge^p(\mathbb{C}^{mn})$ we have:

- if $k < i - 1$ or $k > j + 1$ then $e_k e_\sigma = e_\sigma e_k$.
- if $i \leq k \leq j$ then $e_k e_\sigma = e_\sigma e_k = 0$.
- if $k < i$ or $k > j$ then $f_k e_\sigma = e_\sigma f_k$.

For $i < j$, let $E_{i,j}$ denote the term $[e_i, [e_{i+1}, [\dots [e_{j-1}, e_j]]]$ and $F_{i,j}$ denote $[[[f_j, f_{j-1}], \dots, f_i]]$.

Lemma 5

$$E_{i,j}(v_c) = \begin{cases} (-1)^{|c \cap [i+1, j]|} v_d & \text{if } j+1 \in c \text{ and } i \notin c, \text{ where } d = c - \{j+1\} + \{i\} \\ 0 & \text{otherwise} \end{cases}$$

$$F_{i,j}(v_c) = \begin{cases} (-1)^{|c \cap [i+1, j]|} v_d & \text{if } j+1 \notin c \text{ and } i \in c, \text{ where } d = c - \{i\} + \{j+1\} \\ 0 & \text{otherwise} \end{cases}$$

Proof: We provide a detailed proof for $E_{i,j}$. The proof for $F_{i,j}$ is similar.

We prove this by induction on $j - i$. The base case is when $j - i = 0$. Here, with the convention that $E_{i,i} = e_i$, the lemma follows from the definition of the operator e_i .

For the inductive case (i.e. $i < j$), consider $E_{i,j} = [e_i, E_{i+1,j}] = e_i E_{i+1,j} - E_{i+1,j} e_i$. Thus,

$$E_{i,j}(v_c) = e_i E_{i+1,j}(v_c) - E_{i+1,j} e_i(v_c)$$

Suppose that $E_{i+1,j}(v_c) = 0$, so the first-term in the above expression is zero. Then, by the induction hypothesis, either $j+1 \notin c$ or $i+1 \in c$.

If $i+1 \notin c$, then $j+1 \notin c$. Note that in this case, $e_i(v_c) = 0$. Thus, $E_{i,j}(v_c) = 0$ and $j+1 \notin c$.

If $j+1 \in c$, then $i+1 \in c$. In this case, if $i \in c$, then $e_i(v_c) = 0$ and thus, $E_{i,j}(v_c) = 0$ and $i \in c$. Therefore, we assume that $i \notin c$ alongwith $j+1 \in c$ and $i+1 \in c$. So, we have

$$e_i(v_c) = v_d \text{ where } d = c - \{i+1\} + \{i\}$$

As, $j+1 \in d$ and $i+1 \notin d$, by induction hypothesis,

$$E_{i+1,j}(v_d) = (-1)^{|d \cap [i+2, j]|} v_e \text{ where } e = d - \{j+1\} + \{i+1\}$$

Therefore,

$$\begin{aligned}
E_{i,j}(v_c) &= -E_{i+1,j}e_i(v_c) \\
&= -E_{i+1,j}(v_d) \\
&= -(-1)^{|d \cap [i+2,j]|}v_e \\
&= (-1)^{|c \cap [i+1,j]|}v_e
\end{aligned}$$

The last equation follows from the fact that $i+1 \in c$ and $d = c - \{i+1\} + \{i\}$. Also, observe that $e = c - \{j+1\} + \{i\}$.

Now, we consider the case when $E_{i+1,j}(v_c) \neq 0$. Then, by induction, we have that $j+1 \in c$ and $i+1 \notin c$. Further,

$$E_{i+1,j}(v_c) = (-1)^{|c \cap [i+2,j]|}v_d \text{ where } d = c - \{j+1\} + \{i+1\}$$

Note that, as $i+1 \notin c$, $e_i(v_c) = 0$. Thus, in this case,

$$\begin{aligned}
E_{i,j}(v_c) &= e_i E_{i+1,j}(v_c) - E_{i+1,j}e_i(v_c) \\
&= e_i((-1)^{|c \cap [i+2,j]|}v_d) \\
&= (-1)^{|c \cap [i+2,j]|}e_i(v_d) \\
&= (-1)^{|c \cap [i+1,j]|}e_i(v_d)
\end{aligned}$$

The last equality follows from the observation that $i+1 \notin c$.

If $i \in c$, then $i \in d$ as well and $e_i(v_d) = 0$, consequentially $E_{i,j}(v_c) = 0$ as expected.

If $i \notin c$, then $i \notin d$ as well. As $i+1 \in d$, we have

$$E_{i,j}(v_c) = (-1)^{|c \cap [i+1,j]|}e_i(v_d) = (-1)^{|c \cap [i+1,j]|}v_e$$

where $e = d - \{i+1\} + \{i\} = c - \{j+1\} + \{i\}$.

Q.E.D.

Lemma 6 For i, j, i', j' , on $\wedge^k(\mathbb{C}^{mn})$ we have:

- (i) $[E_{i,j}, E_{i',j'}] = 0$ unless either $j'+1 = i$ or $j+1 = i'$.
- (ii) $[F_{i,j}, E_{i',j'}] = 0$ unless either $j' = j$ or $i' = i$.
- (iii) $E_{i,j}E_{i',j'} = E_{i'j'}E_{ij} = 0$ if $i = i'$ or $j = j'$.
- (iv) $F_{i,j}E_{i',j'} = E_{i'j'}F_{ij} = 0$ if $j+1 = i'$ or $i = j'+1$.

2.2 Commuting actions on $\wedge^k(\mathbb{C}^{mn})$

We are now ready to define two actions, that of $U_q(gl_m)$ and $U_q(gl_n)$ on $\wedge^p(\mathbb{C}^{mn})$. This will consist of some special elements $(E_i^L, F_i^L, q^{\epsilon_i^L})$ and $(E_k^R, F_k^R, q^{\epsilon_k^R})$ which will implement the action of $U_q(sl_m)$ and $U_q(sl_n)$, respectively.

We consider the free $\mathbb{Z}$ -module $\mathbb{E} = \bigoplus_{i=1}^{mn} \mathbb{Z}\epsilon_i$ and define an inner product by extending $\langle \epsilon_i, \epsilon_j \rangle = \delta_{i,j}$. Define $\kappa_{i,j} \in \mathbb{E}$ as $\epsilon_i - \epsilon_j$.

We note that:

Lemma 7 For $\alpha \in \mathbb{E}$, we have:

- $e_j q^\alpha = q^{\langle \alpha, \kappa_{j+1,j} \rangle} q^\alpha e_j$.
- $f_j q^\alpha = q^{\langle \alpha, \kappa_{j,j+1} \rangle} q^\alpha f_j$.
- $E_{i,j} q^\alpha = q^{\langle \alpha, \kappa_{j+1,i} \rangle} q^\alpha E_{i,j}$.

Next, we define the **left operators** using:

$$\begin{aligned} B_i^k &= \sum_{j=0}^{k-2} -h_{jm+i} \\ A_i^k &= \sum_{j=k}^{n-1} h_{jm+i} \end{aligned}$$

We define the map $\phi_L : U_q(gl_m) \rightarrow U_q(gl_{mn})$ as:

$$\begin{aligned} q^{\epsilon_i^L} &= \phi_L(q^{\epsilon_i}) = \prod_{j=0}^{n-1} q^{\epsilon_{jm+i}} \\ E_i^L &= \phi_L(e_i) = e_i + q^{-h_i} e_{m+i} + \dots (\prod_{j=0}^{n-2} q^{-h_{jm+i}}) e_{(n-1)m+i} \\ &= q^{B_i^1} e_i + q^{B_i^2} e_{m+i} + \dots q^{B_i^n} e_{(n-1)m+i} \\ F_i^L &= \phi_L(f_i) = (\prod_{j=1}^{n-1} q^{h_{jm+i}}) f_i + \dots + q^{h_{(n-1)m+i}} f_{(n-2)m+i} + f_{(n-1)m+i} \\ &= q^{A_i^1} f_i + \dots + q^{A_i^{n-1}} f_{(n-2)m+i} + q^{A_i^n} f_{(n-1)m+i} \end{aligned}$$

Proposition 8 *The map $\phi_L : U_q(gl_m) \rightarrow U_q(gl_{mn})$ is an algebra homomorphism.*

Proof: The embedding of $\phi_L : U_q(gl_m) \rightarrow U_q(gl_{mn})$ actually comes from:

$$U_q(gl_m) \xrightarrow{\Delta} U_q(gl_m) \otimes \dots U_q(gl_m) \rightarrow U_q(gl_{mn})$$

where (i) there are n copies in the tensor-product, and (ii) Δ is the n -way Hopf. This verifies that ϕ_L is an algebra map.

We define the **right operators**:

Definition 9

$$\begin{aligned} \beta_i^k &= \sum_{j=i+1}^m \epsilon_{km+j} - \sum_{j=i+1}^m \epsilon_{(k-1)m+j} \\ \alpha_i^k &= \sum_{j=1}^{i-1} \epsilon_{i(k-1)m+j} - \sum_{j=1}^{i-1} \epsilon_{km+j} \end{aligned}$$

We define the “map” $\phi_R : U_q(gl_n) \rightarrow U_q(gl_{mn})$ as:

$$\begin{aligned} \phi_R(q^{\epsilon_k^R}) &= \prod_{i=1}^m q^{\epsilon_{(k-1)m+i}} \\ \phi_R(E_k^R) &= \sum_{i=1}^m q^{\beta_i^k} E_{(k-1)m+i, km+i-1} \\ \phi_R(F_k^R) &= \sum_{i=1}^m q^{\alpha_i^k} F_{(k-1)m+i, km+i-1} \\ \phi_R(h_k^R) &= \sum_{i=1}^m \epsilon_{(k-1)m+i} - \epsilon_{km+i} \end{aligned}$$

Remark: ϕ_R serves merely to identify a set of elements in $U_q(gl_{mn})$ corresponding to the generators of $U_q(gl_m)$. Thus, while $\phi_L : U_q(gl_m) \rightarrow U_q(gl_{mn})$ is an algebra homomorphism, the corresponding statement for $U_q(gl_n)$ is not true. However, as we will show that the composites:

$$\begin{aligned} U_q(gl_m) &\xrightarrow{\phi_L} U_q(gl_{mn}) \longrightarrow \text{End}_{\mathbb{C}(q)}(\wedge^p(\mathbb{C}^{mn})) \\ U_q(gl_n) &\xrightarrow{\phi_R} U_q(gl_{mn}) \longrightarrow \text{End}_{\mathbb{C}(q)}(\wedge^p(\mathbb{C}^{mn})) \end{aligned}$$

are **commuting algebra homomorphisms** making $\wedge^p(\mathbb{C}^{mn})$ into a $U_q(gl_m) \otimes U_q(gl_n)$ -module.

We will identify $\mathbb{C}^{mn}$ as $\mathbb{C}^m \otimes \mathbb{C}^n$ arranging the typical element in an $m \times n$ array, reading column-wise from left to right, and within each column from top to bottom (see below). In this notation, see Fig. 1 for individual terms of the left operators and Fig. 2 for the right operators.

1	6	11	16
2	7	12	17
3	8	13	18
4	9	14	19
5	10	15	20




Figure 1: Terms in the Left Operators




Figure 2: Terms in the Right Operators

2.3 Proofs

For an operator $O = q^\mu E_{i,j}$ (where $\mu \in \mathbb{E}$ is arbitrary) let us define $\kappa(O) = \epsilon_{j+1} - \epsilon_i$ and for the operator $O = q^\mu F_{i,j}$, we define $\kappa(O)$ as $\epsilon_i - \epsilon_{j+1}$. We extend this notation so that $E_{i,i} = e_i$ (with $\kappa(E_{i,i}) = \epsilon_{i+1} - \epsilon_i$) and $F_{j,j} = f_j$ (with $\kappa(F_{j,j}) = \epsilon_j - \epsilon_{j+1}$).

We define $\mathcal{L}$ and $\mathcal{R}$ as two sets of operators:

$$\begin{aligned} \mathcal{L} &= \{q^{B_i^k} e_{(k-1)m+i}, q^{A_i^k} f_{(k-1)m+i} | 1 \leq i \leq m-1, 1 \leq k \leq n\} \\ \mathcal{R} &= \{q^{\beta_i^k} E_{(k-1)m+i, km+i-1}, q^{\alpha_i^k} F_{(k-1)m+i, km+i-1} | 1 \leq i \leq m, 1 \leq k \leq n-1\} \end{aligned}$$

Notice that we may write $E_i^L = \sum_p l_{ip}$ and $E_k^R = \sum_j r_{kj}$ where $l_{ip} \in \mathcal{L}$ and $r_{kj} \in \mathcal{R}$. Whence $[E_i^L, E_k^R]$ is expressible as lie-brackets of elements of $\mathcal{L}$ and $\mathcal{R}$. Of course, we wish to show that $[E_i^L, E_k^R]$ and its three cousins are actually zero.

Lemma 10 *For any $L \in \mathcal{L}$ and any $R \in \mathcal{R}$ if $\langle \kappa(L), \kappa(R) \rangle \geq 0$ then $[L, R] = 0$.*

Proof: We first take the case when $\langle \kappa(L), \kappa(R) \rangle = 0$. We take for example $L = q^{B_{i'}^{k'}} e_{(k'-1)m+i'}$ and $R = q^{\alpha_i^k} F_{(k-1)m+i, km+i-1}$. The condition $\langle \kappa(L), \kappa(R) \rangle = 0$ implies (see Figs. 1, 2) that

$$\begin{aligned} F_{(k-1)m+i, km+i-1} q^{B_{i'}^{k'}} &= q^{B_{i'}^{k'}} F_{(k-1)m+i, km+i-1} \\ e_{(k'-1)m+i'} q^{\alpha_i^k} &= q^{\alpha_i^k} e_{(k'-1)m+i'} \end{aligned}$$

Whence

$$[L, R] = q^{B_{i'}^{k'} + \alpha_i^k} [e_{(k'-1)m+i'}, e_{(k-1)m+i, km+i-1}] = 0$$

where the last equality follows from Lemma 6 (ii).

For the case with $\langle \kappa(L), \kappa(R) \rangle = 1$, Lemma 6, parts (iii), (iv), immediately implies an even stronger claim. Q.E.D.

Thus the only non-commuting (L, R) pairs are shown in Fig. 3.




Figure 3: The Eight Non-Commuting Terms

By lemma 10, for the purpose of showing commutation we may as well assume that $n = m = 2$. The following argument assumes $n = 2$ but retains m for notational convenience. In other words, we have:

$$\begin{aligned} E_i^L &= e_i + q^{-h_i} e_{m+i} \\ F_i^L &= q^{h_i} f_i + f_{m+i} \end{aligned}$$

For $i = 1, \dots, m$ define $\beta_i, \alpha_i \in \mathbb{E}$ as

$$\begin{aligned} \beta_i &= \sum_{j=i+1}^m \epsilon_{m+j} - \sum_{j=i+1}^m \epsilon_j \\ \alpha_i &= \sum_{j=1}^{i-1} \epsilon_j - \sum_{j=1}^{i-1} \epsilon_{m+j} \end{aligned}$$

Next, define

$$\begin{aligned} E^R &= \sum_{i=1}^m q^{\beta_i} E_{i,m+i-1} \\ F^R &= \sum_{i=1}^m q^{\alpha_i} F_{i,m+i-1} \\ h^R &= \sum_{i=1}^m \epsilon_i - \epsilon_{m+i} \end{aligned}$$

Note that $E_1^R = E^R$, $F_1^R = F^R$ and $h_1^R = h^R$.

Lemma 11 For $1 \leq i \leq m-1$,

- $[e_i, q^{\beta_{i+1}} E_{i+1,m+i}] = q^{\beta_{i+1}} E_{i,m+i}$.
- $[q^{-h_i} e_{m+i}, q^{\beta_i} E_{i,m+i-1}] = qq^{\beta_i - h_i} [e_{m+i}, E_{i,m+i-1}]$.

Proof: We prove the first assertion below. We start with analyzing

$$\begin{aligned} [e_i, q^{\beta_{i+1}} E_{i+1,m+i}] &= e_i q^{\beta_{i+1}} E_{i+1,m+i} - q^{\beta_{i+1}} E_{i+1,m+i} e_i \\ &= q^{<\beta_{i+1}, -h_i>} q^{\beta_{i+1}} e_i E_{i+1,m+i} - q^{\beta_{i+1}} E_{i+1,m+i} e_i \end{aligned}$$

A small calculation shows that $<\beta_{i+1}, -h_i> = 0$. Therefore,

$$\begin{aligned} [e_i, q^{\beta_{i+1}} E_{i+1,m+i}] &= q^{\beta_{i+1}} (e_i E_{i+1,m+i} - E_{i+1,m+i} e_i) \\ &= q^{\beta_{i+1}} E_{i,m+i} \end{aligned}$$

Now, we turn to the second claim. Towards this, we expand $[q^{-h_i}e_{m+i}, q^{\beta_i}E_{i,m+i-1}]$ as

$$\begin{aligned} &= q^{-h_i}e_{m+i}q^{\beta_i}E_{i,m+i-1} - q^{\beta_i}E_{i,m+i-1}q^{-h_i}e_{m+i} \\ &= q^{-h_i}q^{<\beta_i, -h_{m+i}>}q^{\beta_i}e_{m+i}E_{i,m+i-1} - q^{\beta_i}q^{<-h_i, \kappa_{m+i,i}>}q^{-h_i}E_{i,m+i-1}e_{m+i} \end{aligned}$$

We observe that $<\beta_i, -h_{m+i}> = 1$ and $<-h_i, \kappa_{m+i,i}> = 1$. Therefore,

$$\begin{aligned} [q^{-h_i}e_{m+i}, q^{\beta_i}E_{i,m+i-1}] &= q^{\beta_i-h_i}(qe_{m+i}E_{i,m+i-1} - qE_{i,m+i-1}e_{m+i}) \\ &= q.q^{\beta_i-h_i}[e_{m+i}, E_{i,m+i-1}] \end{aligned}$$

Q.E.D.

Lemma 12 $[E_i^L, E^R] = 0$

Proof:

$$\begin{aligned} [E_i^L, E^R] &= [e_i + q^{-h_i}e_{m+i}, \sum_{j=1}^m q^{\beta_j}E_{j,m+j-1}] \\ &= [e_i, q^{\beta_{i+1}}E_{i+1,m+i}] + [q^{-h_i}e_{m+i}, q^{\beta_i}E_{i,m+i-1}] \\ &= q^{\beta_{i+1}}E_{i,m+i} + qq^{\beta_i-h_i}(e_{m+i}E_{i,m+i-1} - E_{i,m+i-1}e_{m+i}) \end{aligned}$$

As $\beta_i = \beta_{i+1} + \epsilon_{m+i+1} - \epsilon_{i+1}$, $\beta_i - h_i = \beta_{i+1} + \epsilon_{m+i+1} - \epsilon_i = \beta_{i+1} + \kappa_{m+i+1,i}$.

$$[E_i^L, E^R] = q^{\beta_{i+1}}(E_{i,m+i} + q.q^{\kappa_{m+i+1,i}}(e_{m+i}E_{i,m+i-1} - E_{i,m+i-1}e_{m+i}))$$

Now we evaluate the outer bracket at v_c . So, we are looking at (*)

$$E_{i,m+i}(v_c) + q.q^{\kappa_{m+i+1,i}}(e_{m+i}E_{i,m+i-1}(v_c) - E_{i,m+i-1}e_{m+i}(v_c))$$

If $m+i+1 \notin c$, then all the three terms in the above expression evaluate to 0. The middle term certainly evaluates to 0 after the application of e_{m+i} (even if $E_{i,m+i-1}(v_c) \neq 0$).

Similarly, if $i \in c$, then all the three terms evaluate to 0.

So, henceforth, we work with the assumption that $m+i+1 \in c$ and $i \notin c$.

Now, we consider the case where $m+i \in c$. In this case, with $c_1 = c - \{m+i+1\} + \{i\}$ and $c_2 = c - \{m+i\} + \{i\}$, (*) evaluates to

$$\begin{aligned} * &= (-1)^{|c \cap [i+1, m+i]|}v_{c_1} + q.q^{\kappa_{m+i+1,i}}e_{m+i}((-1)^{|c \cap [i+1, m+i-1]|}v_{c_2}) \\ &= (-1)^{|c \cap [i+1, m+i-1]|}(-v_{c_1} + q.q^{\kappa_{m+i+1,i}}v_{c_1}) \\ &= (-1)^{|c \cap [i+1, m+i-1]|}\left(-v_{c_1} + q.\frac{1}{q}v_{c_1}\right) \\ &= 0 \end{aligned}$$

Now, we consider the remaining case where $m+i \notin c$. In this case, with the notation $c_1 = c - \{m+i+1\} + \{i\}$ and $c_2 = c - \{m+i+1\} + \{m+i\}$, (*) evaluates to

$$\begin{aligned} * &= (-1)^{|c \cap [i+1, m+i]|}v_{c_1} - q.q^{\kappa_{m+i+1,i}}E_{i,m+i-1}(v_{c_2}) \\ &= (-1)^{|c \cap [i+1, m+i]|}v_{c_1} - q.q^{\kappa_{m+i+1,i}}((-1)^{|c_2 \cap [i+1, m+i-1]|}v_{c_1}) \\ &= (-1)^{|c \cap [i+1, m+i-1]|}(v_{c_1} - q.q^{\kappa_{m+i+1,i}}v_{c_1}) \\ &= (-1)^{|c \cap [i+1, m+i-1]|}\left(v_{c_1} - q.\frac{1}{q}v_{c_1}\right) \\ &= 0 \end{aligned}$$

Q.E.D.

Lemma 13 For $1 \leq i \leq m-1$,

- $[f_i q^{h_{m+i}}, q^{\beta_i}E_{i,m+i-1}] = qq^{h_{m+i}+\beta_i}[f_i, E_{i,m+i-1}]$.
- $[f_{m+i}, q^{\beta_{i+1}}E_{i+1,m+i}] = q^{\beta_{i+1}}[f_{m+i}, E_{i+1,m+i}]$.

Proof: We start by proving the first claim.

$$\begin{aligned}
[f_i q^{h_{m+i}}, q^{\beta_i} E_{i,m+i-1}] &= f_i q^{h_{m+i}} q^{\beta_i} E_{i,m+i-1} - q^{\beta_i} E_{i,m+i-1} f_i q^{h_{m+i}} \\
f_i q^{h_{m+i}} q^{\beta_i} E_{i,m+i-1} &= q^{<h_{m+i}+\beta_i, \kappa_{i,i+1}>} q^{h_{m+i}+\beta_i} f_i E_{i,m+i-1} \\
&= q q^{h_{m+i}+\beta_i} f_i E_{i,m+i-1} \\
q^{\beta_i} E_{i,m+i-1} f_i q^{h_{m+i}} &= q^{\beta_i} E_{i,m+i-1} q^{<h_{m+i}, \kappa_{i,i+1}>} q^{h_{m+i}} f_i \\
&= q^{<h_{m+i}, \kappa_{i,i+1}+\kappa_{m+i,i}>} q^{h_{m+i}+\beta_i} E_{i,m+i-1} f_i \\
&= q q^{h_{m+i}+\beta_i} E_{i,m+i-1} f_i
\end{aligned}$$

Thus,

$$\begin{aligned}
[f_i q^{h_{m+i}}, q^{\beta_i} E_{i,m+i-1}] &= q q^{h_{m+i}+\beta_i} (f_i E_{i,m+i-1} - E_{i,m+i-1} f_i) \\
&= q q^{h_{m+i}+\beta_i} [f_i, E_{i,m+i-1}]
\end{aligned}$$

Now, we turn to the second claim.

$$\begin{aligned}
[f_{m+i}, q^{\beta_{i+1}} E_{i+1,m+i}] &= f_{m+i} q^{\beta_{i+1}} E_{i+1,m+i} - q^{\beta_{i+1}} E_{i+1,m+i} f_{m+i} \\
f_{m+i} q^{\beta_{i+1}} E_{i+1,m+i} &= q^{<\beta_{i+1}, \kappa_{m+i}, m+i+1>} q^{\beta_{i+1}} f_{m+i} E_{i+1,m+i} \\
&= q^{\beta_{i+1}} f_{m+i} E_{i+1,m+i}
\end{aligned}$$

Thus,

$$\begin{aligned}
[f_{m+i}, q^{\beta_{i+1}} E_{i+1,m+i}] &= q^{\beta_{i+1}} f_{m+i} E_{i+1,m+i} - q^{\beta_{i+1}} E_{i+1,m+i} f_{m+i} \\
&= q^{\beta_{i+1}} [f_{m+i}, E_{i+1,m+i}]
\end{aligned}$$

Lemma 14 $[F_i^L, E^R] = 0$

Proof:

$$\begin{aligned}
[F_i^L, E^R] &= [f_i q^{h_{m+i}} + f_{m+i}, \sum_{j=1}^m q^{\beta_j} E_{j,m+j-1}] \\
&= [f_i q^{h_{m+i}}, q^{\beta_i} E_{i,m+i-1}] + [f_{m+i}, q^{\beta_{i+1}} E_{i+1,m+i}] \\
&= q q^{h_{m+i}+\beta_i} [f_i, E_{i,m+i-1}] + q^{\beta_{i+1}} [f_{m+i}, E_{i+1,m+i}]
\end{aligned}$$

As $\beta_i = \beta_{i+1} + \epsilon_{m+i+1} - \epsilon_{i+1}$, $\beta_i + h_{m+i} = \beta_{i+1} + \epsilon_{m+i} - \epsilon_{i+1} = \beta_{i+1} + \kappa_{m+i,i+1}$.

$$[E_i^L, E^R] = q^{\beta_{i+1}} (q q^{\kappa_{m+i,i+1}} [f_i, E_{i,m+i-1}] + [f_{m+i}, E_{i+1,m+i}])$$

Now we evaluate the outer bracket at v_c . So, we are looking at (*)

$$q q^{\kappa_{m+i,i+1}} (f_i E_{i,m+i-1}(v_c) - E_{i,m+i-1} f_i(v_c)) + f_{m+i} E_{i+1,m+i}(v_c) - E_{i+1,m+i} f_{m+i}(v_c)$$

If $m+i \notin c$, then all the four terms in the above expression evaluate to 0. Similarly, if $i+1 \in c$, then all the four terms evaluate to 0.

So, henceforth, we work with the assumption that $m+i \in c$ and $i+1 \notin c$.

Now, we consider the case where $i \in c$. In this case, the first term evaluates to 0. If we further assume that $m+i+1 \notin c$, then the third term also evaluates to 0. Overall, with $c_1 = c - \{i\} + \{i+1\}$ and $c_2 = c - \{m+i\} + \{m+i+1\}$, (*) evaluates to

$$* = -q q^{\kappa_{m+i,i+1}} E_{i,m+i-1}(v_{c_1}) - E_{i+1,m+i}(v_{c_2})$$

With the notation $d = c - \{m+i\} + \{i+1\}$,

$$\begin{aligned}
* &= -q q^{\kappa_{m+i,i+1}} (-1)^{|c_1 \cap [i+1, m+i-1]|} v_d - (-1)^{|c_2 \cap [i+2, m+i]|} v_d \\
&= (-1)^{|c \cap [i+2, m+i-1]|} (q q^{\kappa_{m+i,i+1}} v_d - v_d) \\
&= (-1)^{|c \cap [i+2, m+i-1]|} \left(q \frac{1}{q} v_d - v_d \right) \\
&= 0
\end{aligned}$$

Now we work with the assumptions $i \in c$ and $m+i+1 \in c$ and evaluate (*). With these assumptions, the first and the last term of (*) evaluate to 0. Here, with $c_1 = c - \{i\} + \{i+1\}$, $c_2 = c - \{m+i+1\} + \{i+1\}$ and $d = c - \{m+i\} + \{i+1\}$, (*) evaluates to

$$\begin{aligned}
* &= -qq^{\kappa_{m+i,i+1}} E_{i,m+i-1}(v_{c_1}) + f_{m+i}((-1)^{|c \cap [i+2, m+i]|} v_{c_2}) \\
&= -qq^{\kappa_{m+i,i+1}} (-1)^{|c_1 \cap [i+1, m+i-1]|} v_d + (-1)^{|c \cap [i+2, m+i]|} v_d \\
&= (-1)^{|c \cap [i+2, m+i-1]|} (qq^{\kappa_{m+i,i+1}} v_d - v_d) \\
&= (-1)^{|c \cap [i+2, m+i-1]|} \left(q \frac{1}{q} v_d - v_d \right) \\
&= 0
\end{aligned}$$

Now we consider the case with $i \notin c$. In this case, the second term in (*) evaluates to 0. As before, if we further assume that $m+i+1 \notin c$, then the third term also evaluates to 0. Overall, with $c_1 = c - \{m+i\} + \{i\}$, $c_2 = c - \{m+i\} + \{m+i+1\}$, and $d = c - \{m+i\} + \{i+1\}$,

$$\begin{aligned}
* &= qq^{\kappa_{m+i,i+1}} f_i((-1)^{|c \cap [i+1, m+i-1]|} v_{c_1}) - E_{i+1, m+i}(v_{c_2}) \\
&= qq^{\kappa_{m+i,i+1}} (-1)^{|c \cap [i+1, m+i-1]|} v_d - (-1)^{|c_2 \cap [i+2, m+i]|} v_d \\
&= (-1)^{|c \cap [i+2, m+i-1]|} (qq^{\kappa_{m+i,i+1}} v_d - v_d) \\
&= (-1)^{|c \cap [i+2, m+i-1]|} \left(q \frac{1}{q} v_d - v_d \right) \\
&= 0
\end{aligned}$$

For the only remaining case, we have the assumptions $i \notin c$ and $m+i+1 \in c$. Here, with $c_1 = c - \{m+i\} + \{i\}$, $c_2 = c - \{m+i+1\} + \{i+1\}$ and $d = c - \{m+i\} + \{i+1\}$, (*) evaluates to

$$\begin{aligned}
* &= qq^{\kappa_{m+i,i+1}} f_i((-1)^{|c \cap [i+1, m+i-1]|} v_{c_1}) + f_{m+i}((-1)^{|c \cap [i+2, m+i]|} v_{c_2}) \\
&= qq^{\kappa_{m+i,i+1}} (-1)^{|c \cap [i+1, m+i-1]|} v_d + (-1)^{|c \cap [i+2, m+i]|} v_d \\
&= (-1)^{|c \cap [i+2, m+i-1]|} (qq^{\kappa_{m+i,i+1}} v_d - v_d) \\
&= (-1)^{|c \cap [i+2, m+i-1]|} \left(q \frac{1}{q} v_d - v_d \right) \\
&= 0
\end{aligned}$$

Q.E.D.

We have shown that $[E_i^L, E^R] = 0$ and $[F_i^L, E^R] = 0$. One can similarly show that $[E_i^L, F^R] = [F_i^L, F^R] = 0$. We now prepare towards proving $[F_R, E_R] = (q^{-h^R} - q^{h^R})/(q - q^{-1})$.

Lemma 15 *For $i \neq j$, we have:*

$$[q^{\alpha_i} F_{i,m+i-1}, q^{\beta_j} E_{j,m+j-1}] = 0$$

Proof:

$$\begin{aligned}
[q^{\alpha_i} F_{i,m+i-1}, q^{\beta_j} E_{j,m+j-1}] &= q^{\alpha_i} F_{i,m+i-1} q^{\beta_j} E_{j,m+j-1} - q^{\beta_j} E_{j,m+j-1} q^{\alpha_i} F_{i,m+i-1} \\
&= q^{\alpha_i + \beta_j} (q^{\beta_j(i) - \beta_j(m+i)} F_{i,m+i-1} E_{j,m+j-1} - q^{\alpha_i(m+j) - \alpha_i(j)} E_{j,m+j-1} F_{i,m+i-1}) \\
&= q^a q^{\alpha_i + \beta_j} [F_{i,m+i-1}, E_{j,m+j-1}]
\end{aligned}$$

for an appropriate integer a depending on the whether $i \leq j$ or not. Now, the only material case for v_c is when $i, m+j \in c$ and $j, m+i \notin c$. We may then verify that $[F_{i,m+i-1}, E_{j,m+j-1}] v_c = 0$. Q.E.D.

Lemma 16 *For $1 \leq i \leq m$*

$$[q^{\alpha_i} F_{i,m+i-1}, q^{\beta_i} E_{i,m+i-1}] = q^{\alpha_i + \beta_i} [F_{i,m+i-1}, E_{i,m+i-1}]$$

$$\begin{aligned}
[q^{\alpha_i} F_{i,m+i-1}, q^{\beta_i} E_{i,m+i-1}] &= q^{\alpha_i} F_{i,m+i-1} q^{\beta_i} E_{i,m+i-1} - q^{\beta_i} E_{i,m+i-1} q^{\alpha_i} F_{i,m+i-1} \\
&= q^{\alpha_i + \beta_i} (q^{\beta_i(i) - \beta_i(m+i)} F_{i,m+i-1} E_{i,m+i-1} - q^{\alpha_i(m+i) - \alpha_i(i)} E_{i,m+i-1} F_{i,m+i-1}) \\
&= q^{\alpha_i + \beta_i} [F_{i,m+i-1}, E_{i,m+i-1}]
\end{aligned}$$

This proves the lemma. Q.E.D.

Define $\delta_j = \epsilon_j - \epsilon_{m+j}$ and let $v_c \in \wedge^p(\mathbb{C}^{mn})$.

Lemma 17

$$(q - q^{-1})[F_{i,m+i-1}, E_{i,m+i-1}]v_c = (q^{-\delta_i} - q^{\delta_i})v_c$$

Proof: If both $i, m+i \in c$ or both $i, m+i \notin c$ then the equality clearly holds. Now if $i \in c, m+i \notin c$ then $q^{\delta_i}v_c = qv_c$ and we have:

$$\begin{aligned} (q - q^{-1})[F_{i,m+i-1}, E_{i,m+i-1}]v_c &= (q - q^{-1})(-v_c) \\ &= (q^{-\delta_i} - q^{\delta_i})v_c \end{aligned}$$

On the other hand, if $i \notin c, m+i \in c$, then $q^{\delta_i}v_c = q^{-1}v_c$ and we have:

$$\begin{aligned} (q - q^{-1})[F_{i,m+i-1}, E_{i,m+i-1}]v_c &= (q - q^{-1})(v_c) \\ &= (q^{-\delta_i} - q^{\delta_i})v_c \end{aligned}$$

This proves the lemma.

We now prove:

Proposition 18 Let $h_R = \sum_{i=1}^m \epsilon_i - \epsilon_{m+i}$ then

$$[F^R, E^R] = \frac{q^{-h_R} - q^{h_R}}{q - q^{-1}}$$

Proof: By the above lemmas, we have:

$$[F^R, E^R] = \sum_{i=1}^m q^{\alpha_i + \beta_i} [F_{i,m+i-1}, E_{i,m+i-1}]$$

Whence

$$\begin{aligned} (q - q^{-1})[F^R, E^R]v_c &= \sum_{i=1}^m (q^{-\delta_i} - q^{\delta_i})q^{\alpha_i + \beta_i}v_c \\ &= \sum_{i=1}^m q^{\alpha_i + \beta_i - \delta_i}v_c - q^{\alpha_i + \beta_i + \delta_i}v_c \end{aligned}$$

Now

$$\alpha_i + \beta_i = \left(\sum_{j=1}^{i-1} \delta_j \right) - \left(\sum_{j=i+1}^m \delta_j \right)$$

and thus

$$\alpha_i + \beta_i - \delta_i = \alpha_{i-1} + \beta_{i-1} + \delta_{i-1} = \left(\sum_{j=1}^{i-1} \delta_j \right) - \left(\sum_{j=i}^m \delta_j \right)$$

Consequently

$$\begin{aligned} (q - q^{-1})[F^R, E^R]v_c &= \sum_{i=1}^m (q^{-\delta_i} - q^{\delta_i})q^{\alpha_i + \beta_i}v_c \\ &= \sum_{i=1}^m q^{\alpha_i + \beta_i - \delta_i}v_c - q^{\alpha_i + \beta_i + \delta_i}v_c \\ &= (q^{\alpha_1 + \beta_1 - \delta_1} - q^{\alpha_m + \beta_m + \delta_m})v_c \\ &= (q^{-h_R} - q^{h_R})v_c \end{aligned}$$

This proves the proposition. Q.E.D.

We next prove the braid identity.

Definition 19 For $i = 1, \dots, m$ define $\beta_i, \alpha_i \in \mathbb{E}$ as

$$\begin{aligned} \beta_i &= \sum_{j=i+1}^m \epsilon_{m+j} - \sum_{j=i+1}^m \epsilon_j \\ \beta_i^* &= \sum_{j=i+1}^m \epsilon_{2m+j} - \sum_{j=i+1}^m \epsilon_{m+j} \end{aligned}$$

Next, define

$$\begin{aligned} E^R &= \sum_{i=1}^m q^{\beta_i} E_{i,m+i-1} \\ E^{*R} &= \sum_{i=1}^m q^{\beta_i^*} E_{m+i, 2m+i-1} \end{aligned}$$

Note that $E^{*R} = E_2^R$. We will show that:

$$(E^R)^2 E^{*R} - (q + q^{-1}) E^R E^{*R} E^R + E^{*R} (E^R)^2 = 0$$

We define $g_i = q^{\beta_i} E_{i,m+i-1}$ and $g_j^* = q^{\beta_j^*} E_{m+j,2m+j-1}$.

Lemma 20 *For distinct $i, j, k \in [m]$ and on $\wedge^p(\mathbb{C}^{mn})$, we have that*

$$(g_i g_j + g_j g_i) g_k^* - (q + q^{-1})(g_i g_k^* g_j + g_j g_k^* g_i) + g_k^*(g_i g_j + g_j g_i) = 0$$

Proof: Let us prove this in several cases. In all cases, we will use:

$$\begin{aligned} E_{i,m+i-1} q^{\beta_j} &= \begin{cases} q^2 q^{\beta_j} E_{i,m+i-1} & \text{if } i > j \\ q^{\beta_j} E_{i,m+i-1} & \text{if } i \leq j \end{cases} \\ E_{i,m+i-1} q^{\beta_j^*} &= \begin{cases} q^{-1} q^{\beta_j^*} E_{i,m+i-1} & \text{if } i > j \\ q^{\beta_j^*} E_{i,m+i-1} & \text{if } i \leq j \end{cases} \\ E_{m+i,2m+i-1} q^{\beta_j} &= \begin{cases} q^{-1} q^{\beta_j} E_{i,m+i-1} & \text{if } i > j \\ q^{\beta_j} E_{i,m+i-1} & \text{if } i \leq j \end{cases} \end{aligned}$$

We first consider the case $i < j < k$ and v_c such that $v = E_{i,m+i-1} E_{j,j+m-1} E_{m+k,2m+k-1} v_c$, where, by Lemma 6, the sequence of the operators does not matter. Note further that $g_i g_j g_k^*(v_c) = v^* = q^a \cdot v$. We suppress the factor q^a uniformly in this proof and in the next lemma as well. We see that:

$$\begin{aligned} (g_i g_j + g_j g_i) g_k^* v_c &= (1 + q^2) E_{i,m+i-1} E_{j,j+m-1} E_{m+k,2m+k-1} v_c \\ &= (1 + q^2) v \\ g_k^*(g_i g_j + g_j g_i) v_c &= (q^{-2} + q^2 \cdot q^{-2}) E_{i,m+i-1} E_{j,j+m-1} E_{m+k,2m+k-1} v_c \\ &= (q^{-2} + 1) v \\ (g_i g_k^* g_j + g_j g_k^* g_i) v_c &= (q^{-1} + q^{-1} \cdot q^2) E_{i,m+i-1} E_{j,j+m-1} E_{m+k,2m+k-1} v_c \\ &= (q^{-1} + q) v \end{aligned}$$

This proves the assertion for $i < j < k$.

Next, let us consider $i < k < j$:

$$\begin{aligned} (g_i g_j + g_j g_i) g_k^* v_c &= (q^{-1} + q) v \\ g_k^*(g_i g_j + g_j g_i) v_c &= (q^{-1} + q) v \\ (g_i g_k^* g_j + g_j g_k^* g_i) v_c &= 2v \end{aligned}$$

This proves the assertion for $i < k < j$.

Next, let us consider $k < i < j$:

$$\begin{aligned} (g_i g_j + g_j g_i) g_k^* v_c &= (1 + q^{-2}) v \\ g_k^*(g_i g_j + g_j g_i) v_c &= (1 + q^2) v \\ (g_i g_k^* g_j + g_j g_k^* g_i) v_c &= (q + q^{-1}) v \end{aligned}$$

This proves the assertion for $k < i < j$ and completes the proof of the lemma. Q.E.D.

Lemma 21 *For distinct $i, j \in [m]$ and on $\wedge^p(\mathbb{C}^{mn})$, we have that*

$$(g_i g_j + g_j g_i) g_i^* - (q + q^{-1})(g_i g_i^* g_j + g_j g_i^* g_i) + g_i^*(g_i g_j + g_j g_i) = 0$$

Proof: There are two cases to consider, viz., $g_i g_i^* v_c = 0$ and $g_i^* g_i v_c = 0$. Let us consider the first case, i.e., $g_i g_i^* v_c = 0$, in which case we need to show:

$$-(q + q^{-1}) g_j g_i^* g_i + g_i^*(g_i g_j + g_j g_i) = 0$$

Let v be such that $E_{m+i,2m+i-1}E_{i,m+i-1}E_{j,m+j-1}v_c = v$ (see comment in proof of Lemma 20). We see that for $j > i$:

$$\begin{aligned} g_i^*(g_i g_j + g_j g_i)v_c &= (1 + q^2)v \\ g_j g_i^* g_i v_c &= qv \end{aligned}$$

This proves the lemma for $j > i$. Next, for $j < i$, with $v = E_{j,m+j-1}E_{m+i,2m+i-1}E_{i,m+i-1}v_c$ and we have:

$$\begin{aligned} g_i^*(g_i g_j + g_j g_i)v_c &= (q + q^{-1})v \\ g_j g_i^* g_i v_c &= v \end{aligned}$$

This proves the case when $g_i g_i^* v_c = 0$. The other case is similarly proved. Q.E.D.

Proposition 22 For $E^R = E_1^R$ and $E^{*R} = E_2^R$, we have:

$$(E^R)^2 E^{*R} - (q + q^{-1}) E^R E^{*R} E^R + E^{*R} (E^R)^2 = 0$$

Proof: Let

$$B = (E^R)^2 E^{*R} - (q + q^{-1}) E^R E^{*R} E^R + E^{*R} (E^R)^2$$

For a given v_c , we look at $B \cdot v_c$ and classify the result by the $U_q(gl_{mn})$ weight. We see that the allowed weights are $wt(v_c) - \kappa_{m+i,i} - \kappa_{m+j,j} - \kappa_{m+k,k}$ for various i, j, k . Further, we see that:

$$\begin{aligned} E^R &= \sum_{i=1}^m g_i \\ E^{*R} &= \sum_{i=1}^m g_i^* \end{aligned}$$

is a separation of E^R and E^{*R} by $U_q(gl_{mn})$ -weights. Therefore showing $B \cdot v_c = 0$ amounts to various cases on i, j, k . The main cases are settled by Lemmas 20, 21. Other cases are easier. Q.E.D.

Proposition 23 The map $\phi_R : U_q(gl_n) \rightarrow \text{End}_{\mathbb{C}(q)}(\wedge^p(\mathbb{C}^{mn}))$ is an algebra homomorphism. At $q = 1$, ϕ_R factorizes through $U_q(gl_{mn})$, i.e.,

$$\phi_R(1) : U_1(gl_n) \rightarrow U_1(gl_{mn}) \rightarrow \text{End}_{\mathbb{C}}(\wedge^p(\mathbb{C}^{mn}))$$

The proof is obvious. The family $\{E_k^R, F_k^R, q^{\epsilon_k^R}\}$ satisfy all the properties for $U_q(gl_n)$. Also note that at $q = 1$, $\phi_R(1)$ reduces to the standard injection which commutes with $\phi_L(1)$.

3 The module V_λ

We have thus seen the algebra maps $\phi_L : U_q(gl_m) \rightarrow U_q(gl_{mn}) \rightarrow \text{End}_{\mathbb{C}(q)}(\wedge^k(\mathbb{C}^{mn}))$ and $\phi_R : U_q(gl_n) \rightarrow \text{End}_{\mathbb{C}(q)}(\wedge^k(\mathbb{C}^{mn}))$. Since the two actions commute, this converts $\wedge^k(\mathbb{C}^{mn})$ into a $U_q(gl_m) \otimes U_q(gl_n)$ -module. Also note that at $q = 1$, we have the factorization:

$$\begin{aligned} \phi_L(1) : U_1(gl_m) &\rightarrow U_1(gl_{mn}) \rightarrow \text{End}_{\mathbb{C}}(\wedge^k(\mathbb{C}^{mn})) \\ \phi_R(1) : U_1(gl_n) &\rightarrow U_1(gl_{mn}) \rightarrow \text{End}_{\mathbb{C}}(\wedge^k(\mathbb{C}^{mn})) \end{aligned}$$

Proposition 24 The actions ϕ_L, ϕ_R convert $\wedge^k(\mathbb{C}^{mn})$ into a $U_q(gl_m) \otimes U_q(gl_n)$ module. Furthermore, at $q = 1$ this matches the restriction of the $U_1(gl_{mn})$ action on $\wedge^k(\mathbb{C}^{mn})$ to $U_1(gl_m) \otimes U_1(gl_n)$.

Since, both $U_q(gl_m)$ and $U_q(gl_n)$ are Hopf-algebras, we see that if M, N are $U_q(gl_m) \otimes U_q(gl_n)$ -modules then so is $M \otimes N$. The action of $U_q(gl_m)$ on $M \otimes N$ defined by

$$\Phi_L : U_q(gl_m) \xrightarrow{\Delta} U_q(gl_m) \otimes U_q(gl_m) \xrightarrow{\phi_L \otimes \phi_L} U_q(gl_{mn}) \otimes U_q(gl_{mn}) \rightarrow \text{End}_{\mathbb{C}(q)}(M \otimes N)$$

In the case M, N are $U_q(gl_{mn})$ -modules, we also have:

$$\Phi'_L : U_q(gl_m) \xrightarrow{\phi_L} U_q(gl_{mn}) \xrightarrow{\Delta} U_q(gl_{mn}) \otimes U_q(gl_{mn}) \rightarrow \text{End}_{\mathbb{C}(q)}(M \otimes N)$$

We may similarly define Φ_R

$$\Phi_R : U_q(gl_n) \xrightarrow{\Delta} U_q(gl_n) \otimes U_q(gl_n) \xrightarrow{\phi_R \otimes \phi_R} \text{End}_{\mathbb{C}(q)}(M \otimes N)$$

Again, if M, N are $U_q(gl_{mn})$ -modules, we have at $q = 1$:

$$\Phi'_R(1) : U_1(gl_n) \xrightarrow{\phi_L(1)} U_1(gl_{mn}) \xrightarrow{\Delta} U_1(gl_{mn}) \otimes U_1(gl_{mn}) \rightarrow \text{End}_{\mathbb{C}}(M \otimes N)$$

Proposition 25

- If M, N are $U_q(gl_m) \otimes U_q(gl_n)$ -modules then so is $M \otimes N$, interpreted as $U_q(gl_m) \otimes U_q(gl_n)$ module through Φ_L and Φ_R .
- The maps $\Phi_L = \Phi'_L$ and $\Phi_R = \Phi'_R$ when $q = 1$. Thus Φ_L and Φ_R are deformations of the action of $U_1(gl_{mn})$ restricted to $U_1(gl_m) \otimes U_1(gl_n)$.

The proof of the first part is obvious. For the second part notice that for $q = 1$ both ϕ_L and ϕ_R match the classical injections (algebra homomorphisms) of $U_q(gl_m)$ (or $U_q(gl_n)$) into $U_q(gl_{mn})$.

Unless otherwise stated, for $U_q(gl_{mn})$ -modules M, N , the $U_q(gl_m)$ and $U_q(gl_n)$ structure on $M \otimes N$ will be that arising from Φ_L and Φ_R .

Lemma 26 For the module $\wedge^k(\mathbb{C}^{mn})$ as a $U_q(gl_m) \otimes U_q(gl_n)$ -module, we have:

$$\wedge^k(\mathbb{C}^{mn}) = \sum_{\lambda} V_{\lambda}(\mathbb{C}^m) \otimes V_{\lambda'}(\mathbb{C}^n)$$

where $|\lambda| = k$.

The proof is clear by setting $q = 1$. Q.E.D.

Next, for a $U_1(gl_{mn})$ -module V_{λ} and the standard embedding $U_1(gl_m) \otimes U_1(gl_n)$, let

$$V_{\lambda}(\mathbb{C}^{mn}) = \oplus_{\alpha, \beta} n_{\alpha, \beta}^{\lambda} V_{\alpha}(\mathbb{C}^m) \otimes V_{\beta}(\mathbb{C}^n)$$

Lemma 27 For $a, b \in \mathbb{Z}$, consider $\wedge^{a+1}(\mathbb{C}^{mn}) \otimes \wedge^{b-1}(\mathbb{C}^{mn})$ and $\wedge^a(\mathbb{C}^{mn}) \otimes \wedge^b(\mathbb{C}^{mn})$ as $U_q(gl_m) \otimes U_q(gl_n)$ -modules. Then there exists an $U_q(gl_m) \otimes U_q(gl_n)$ -equivariant injection $\psi_{a,b}$:

$$\psi_{a,b} : \wedge^{a+1}(\mathbb{C}^{mn}) \otimes \wedge^{b-1}(\mathbb{C}^{mn}) \rightarrow \wedge^a(\mathbb{C}^{mn}) \otimes \wedge^b(\mathbb{C}^{mn})$$

If λ is the shape of two columns sized a and b then the co-kernel $\text{cok}(\psi_{a,b})$ may be written as:

$$\text{cok}(\psi_{a,b}) = \oplus_{\alpha, \beta} n_{\alpha, \beta}^{\lambda} V_{\alpha}(\mathbb{C}^m) \otimes V_{\beta}(\mathbb{C}^n)$$

Proof: For $q = 1$ the above map is a classical construction (see, e.g., [5]). This implies that for general q , the multiplicity of the $U_q(gl_m) \otimes U_q(gl_n)$ -module $V_{\alpha}(\mathbb{C}^m) \otimes V_{\beta}(\mathbb{C}^n)$ in $\wedge^{a+1}(\mathbb{C}^{mn}) \otimes \wedge^{b-1}(\mathbb{C}^{mn})$ does not exceed that in $\wedge^a(\mathbb{C}^{mn}) \otimes \wedge^b(\mathbb{C}^{mn})$. Whence a suitable $\psi_{a,b}$ may be constructed respecting the isotypical components of both modules. The second assertion now follows. Q.E.D.

We now proceed to construct the $U_q(gl_m) \otimes U_q(gl_n)$ module W_{λ} . Let $\lambda' = [\mu_1, \dots, \mu_k]$, i.e., λ has k -columns of length $\mu_1, \dots, \mu_k$. Let C^k be the collection of all columns of size k with strictly increasing entries from the set $[mn]$. For $a \geq b$ and $c \in C^a$ and $c' \in C^b$, we say that $c \leq c'$ if for all $1 \leq i \leq b$, we have $c(i) \leq c'(i)$. A basis for W_{λ} will be the set $SS(\lambda, mn)$, i.e., semi-standard tableau of shape λ with entries in $[mn]$. We interpret this basis as $X^{\lambda} \subseteq Z^{\lambda} = \prod_i C^{\mu_i}$. In other words,

$$X^{\lambda} = \{[c_1, \dots, c_k] | c_i \in C^{\mu_i}, c_i \leq c_{i+1}\}$$

We call X^{λ} as **standard** and $Y^{\lambda} = Z^{\lambda} - X^{\lambda}$ as non-standard. We represent $\wedge^p(\mathbb{C}^{mn})$ as in [8], with the basis C^p and construct $M = \otimes_i \wedge^{\mu_i}(\mathbb{C}^{mn})$ with the basis Z^{λ} . Note that M is a $U_q(gl_m) \otimes U_q(gl_n)$ -module.

Lemma 28 *There is a $U_q(\mathfrak{gl}_m) \otimes U_q(\mathfrak{gl}_n)$ -submodule $N \subseteq M$ such that*

- $\dim(N) = |Y^\lambda| = d$ and
- if $b_i = \sum_{t \in Z^\lambda} \gamma_i^t \cdot t$ (for $i = 1, \dots, d$) is a basis for N then the $d \times d$ -matrix $D = (\gamma_i^t)_{i=1, \dots, d}^{t \in Y^\lambda}$ is invertible.

Proof: This again reduces to a choice of $\psi_{a,b}$ for various a, b . We know that for $q = 1$, the above lemma follows from the straightening relations imposed by the maps $\psi_{a,b}$. Whence, for general q , there must exist an open set of such maps $\psi_{a,b}$.

Proposition 29 *There is a $U_q(\mathfrak{gl}_m) \otimes U_q(\mathfrak{gl}_n)$ -module W_λ and a basis w_t for $t \in X^\lambda$ such that:*

- For $q = 1$, the module is isomorphic to $V_\lambda(\mathbb{C}^{mn})$ treated as a $U_1(\mathfrak{gl}_m) \otimes U_1(\mathfrak{gl}_n)$ -module with the vectors w_t as $U_1(\mathfrak{gl}_{mn})$ weight vectors.
- For general q , w_t continue to be $U_q(\mathfrak{gl}_m) \otimes U_q(\mathfrak{gl}_n)$ weight vectors.

Proof: The desired module is M/N .

4 The construction of $\psi_{a,b}$

The structure of W_λ depends intrinsically on the “straightening relations” $\psi_{a,b}$ (for various a, b) of Lemma 27. In this section we will construct a family of maps:

$$\psi_{a,b} : \wedge^{a+1} \otimes \wedge^{b-1} \rightarrow \wedge^a \otimes \wedge^b$$

These maps will have the following important properties:

- $\psi_{a,b}$ will be $U_q(\mathfrak{gl}_m) \otimes U_q(\mathfrak{gl}_n)$ -equivariant, and
- at $q = 1$, they will also be $U_1(\mathfrak{gl}_{mn})$ -equivariant and will match the standard resolution.

This is done in three steps:

- First, the construction of equivariant maps $\psi_a : \wedge^{a+1} \rightarrow \wedge^a \otimes \wedge^1$ and $\psi'_a : \wedge^{a+1} \rightarrow \wedge^1 \otimes \wedge^a$.
- Next, for a module map $\mu : A \rightarrow B$, the construction of the “adjoint” $\mu^* : B \rightarrow A$.
- Finally constructing $\psi_{a,b}$ using ψ_a and ψ_b^* .

We first begin with the adjoint.

4.1 Normal bases

Let us fix the basis $B = \{v_c | c \subseteq [mn], |c| = k\}$ as the basis of $\wedge^k(\mathbb{C}^{mn})$. We define an inner product on $\wedge^k(\mathbb{C}^{mn})$ as follows. For elements $v_c, v_{c'} \in \wedge^k(\mathbb{C}^{mn})$, let $\langle v_c, v_{c'} \rangle = \delta_{c,c'}$. In other words, the inner product is chosen so that B are ortho-normal.

Abusing notation slightly, we denote, for example by $\langle E_i^L c, c' \rangle$ as short-form for $\langle E_i^L(v_c), v_{c'} \rangle$. We have the **EF-Lemma**:

Lemma 30 *For the action of $U_q(\mathfrak{gl}_m)$ and $U_q(\mathfrak{gl}_n)$ as above, on $\wedge^k(\mathbb{C}^{mn})$ as above, we have:*

$$\begin{aligned} q^{-1} q^{h_i^L}(v_{c'}) \langle E_i^L c, c' \rangle &= q q^{h_i^L}(v_c) \langle E_i^L c, c' \rangle = \langle F_i^L c', c \rangle \\ q^{-1} q^{h_i^R}(v_{c'}) \langle E_i^R c, c' \rangle &= q q^{h_i^R}(v_c) \langle E_i^R c, c' \rangle = \langle F_i^R c', c \rangle \end{aligned}$$

Proof: We have:

$$E_i^L(v_c) = (e_i + q^{-h_i}e_{m+i} + \dots (\prod_{j=0}^{n-2} q^{-h_{jm+i}})e_{(n-1)m+i})v_c$$

Now, by examining the gl_{mn} -weights of c, c' , exactly one of these terms will lead to $v_{c'}$, and so

$$\langle E_i^L c, c' \rangle v_{c'} = (\prod_{j=0}^k q^{-h_{jm+i}})e_{((k+1)m+i)}v_c = (\prod_{j=0}^k q^{-h_{jm+i}}(v_{c'})) \cdot v_{c'}$$

Now, we see that:

$$F_i^L(v_c) = ((\prod_{j=1}^{n-1} q^{h_{jm+i}})f_i + \dots + q^{h_{(n-1)m+i}}f_{(n-2)m+i} + f_{(n-1)m+i})v_c$$

It must be the $f_{(k+1)m+i}$ term that led to v_c . Whence, we have:

$$\langle F_i^L c', c \rangle = ((\prod_{j=k+2}^{n-1} q^{h_{jm+i}}) \cdot f_{(k+1)m+i})v_{c'} = (\prod_{j=k+2}^{n-1} q^{h_{jm+i}})v_c$$

But since c, c' differ only in the entry $(k+1)m+i$, we have

- $q^{h_{(k+1)m+i}}(v_c) = q^{-1}$ and $q^{h_{(k+1)m+i}}(v_{c'}) = q$.
- $(\prod_{j=0}^k q^{-h_{jm+i}}(v_{c'})) = (\prod_{j=0}^k q^{-h_{jm+i}}(v_c))$
- $q^{h_i^L}(v_c) = \prod_{j=0}^{n-1} q^{h_{jm+i}}v_c$

Finally,

$$\begin{aligned} qq^{h_i^L}(v_c)\langle E_i^L c, c' \rangle &= q \prod_{j=k+1}^{n-1} q^{h_{jm+i}}(v_c) \\ &= qq^{h_{(k+1)m+i}}(v_c) \prod_{j=k+2}^{n-1} q^{h_{jm+i}}(v_c) \\ &= \langle F_i^L c', c \rangle \end{aligned}$$

Other assertions are similarly proved. Q.E.D.

Definition 31 Let A be a $U_q(gl_m) \otimes U_q(gl_n)$ module, and let $\mathcal{A} = \{a_1, \dots, a_r\}$ be a basis of A of weight vectors. Define an inner product $\langle \cdot, \cdot \rangle$ on A making $\mathcal{A}$ orthogonal. We say that $\mathcal{A}$ is normal if the EF-lemma Lemma 30 holds (with $a, a' \in \mathcal{A}$ replacing c, c').

Lemma 32 Let A, B be $U_q(gl_m) \otimes U_q(gl_n)$ -modules such that $\mathcal{A} = \{a_1, \dots, a_r\}$ and $\mathcal{B} = \{b_1, \dots, b_s\}$ are normal bases for A and B respectively. Then $\mathcal{A} \otimes \mathcal{B}$ is a normal basis for $A \otimes B$ with the inner product $\langle a \otimes b, a' \otimes b' \rangle = \delta_{a \otimes b, a' \otimes b'}$.

Proof: Let consider the element $a \otimes b$, and the elements $a' \otimes b$ and $a \otimes b'$ such that a' appears in $E_i^L a$ and b' appears in $E_i^L b$.

We see that:

$$\begin{aligned} q \cdot q^{h_i^L}(a \otimes b) \langle E_i^L(a \otimes b), a' \otimes b \rangle &= q \cdot q^{h_i^L}(a \otimes b) \langle (E_i^L \otimes 1 + q^{-h_i^L} \otimes E_i^L)(a \otimes b), a' \otimes b \rangle \\ &= q \cdot q^{h_i^L}(a \otimes b) \langle (E_i^L \otimes 1)(a \otimes b), a' \otimes b \rangle \\ &= q \cdot q^{h_i^L}(a \otimes b) \langle E_i^L a, a' \rangle \\ &= q^{h_i^L}(b) [q \cdot q^{h_i^L}(a) \langle E_i^L a, a' \rangle] \\ &= q^{h_i^L}(b) \langle F_i^L a', a \rangle \end{aligned}$$

On the other hand, we have:

$$\begin{aligned}
\langle F_i^L(a' \otimes b), a \otimes b \rangle &= \langle (F_i^L \otimes q^{h_i^L} + 1 \otimes F_i^L)(a' \otimes b), a \otimes b \rangle \\
&= \langle (F_i^L \otimes q^{h_i^L})(a' \otimes b), a \otimes b \rangle \\
&= q^{h_i^L}(b) \langle F_i^L a', a \rangle
\end{aligned}$$

Other cases are similar. Q.E.D.

Let Ξ be the $\mathbb{Z}$ -submodule generated by ϵ_i^L and ϵ_j^R . Let χ be a Ξ -weight and let $\chi' = \chi + h_i^L$. For a module A with a normal base $\mathcal{A}$, let A_χ be the weight-space of weight χ . We see that $E_i^L : A_\chi \rightarrow A_{\chi'}$, while $F_i^L : A_{\chi'} \rightarrow A_\chi$. Let a_χ be the column-vector of elements of $\mathcal{A}$ of weight χ . Let us define matrices E^A, F^A as:

$$E^A a_{\chi'} = E_i^L a_\chi \quad F^A a_\chi = F_i^L a_{\chi'}$$

By the EF-lemma (i.e., Lemma 30),

$$q \cdot q^{<\chi, h_i^L>} E^A = (F^A)^T$$

Now, let A and B be $U_q(gl_m) \otimes U_q(gl_n)$ with normal bases $\mathcal{A}$ and $\mathcal{B}$ respectively. Let $\mu : A \rightarrow B$ be an equivariant map and let μ_χ be a matrix such that:

$$\mu a_\chi = \mu_\chi b_\chi$$

Equivariance implies:

$$\begin{aligned}
\mu \cdot E_i^L a_\chi &= \mu \cdot E^A a_{\chi'} = E^A \mu_\chi b_{\chi'} \\
E_i^L \cdot \mu a_\chi &= E_i^L \mu_\chi b_\chi = \mu_{\chi'} E^B b_{\chi'}
\end{aligned}$$

Or in other words,

$$E^A \mu_\chi = \mu_{\chi'} E^B \quad F^A \mu_{\chi'} = \mu_\chi F^B$$

Transposing the second equivariance condition, we get:

$$(F^A \mu_{\chi'})^T = (\mu_\chi F^B)^T$$

We may simplify this as:

$$\mu_{\chi'}^T (F^A)^T = (F^B)^T \mu_\chi^T$$

and further:

$$q \cdot q^{<\chi, h_i^L>} \mu_{\chi'}^T E_A = q \cdot q^{<\chi, h_i^L>} E_B \mu_\chi^T$$

i.e., finally:

$$\mu_{\chi'}^T E_A = E_B \mu_\chi^T$$

We may similarly prove that

$$\mu_\chi^T F_A = F_B \mu_{\chi'}^T$$

Both these observations immediately imply:

Proposition 33 *Let $\mu : A \rightarrow B$ be an equivariant map, and let μ_χ be defined as above. We construct the map $\mu^* : B \rightarrow A$ as follows. Define μ^* such that:*

$$\mu^* b_\chi = \mu_\chi^T a_\chi$$

Then $\mu^ : B \rightarrow A$ is equivariant.*

4.2 The Construction of ψ_a

In this section we construct the $U_q(gl_m) \otimes U_q(gl_n)$ -equivariant maps

$$\begin{aligned}\psi_a &: \wedge^{a+1} \rightarrow \wedge^a \otimes \wedge^1 \\ \psi'_a &: \wedge^{a+1} \rightarrow \wedge^1 \otimes \wedge^a\end{aligned}$$

Note that $\wedge^1 = \mathbb{C}^{mn} = \mathbb{C}^m \otimes \mathbb{C}^n$. For convenience, we identify $[mn]$ with $[m] \times [n]$. Under this identification, an element $(i, j) \in [m] \times [n]$ maps to the element $m * (j - 1) + i$.

In this notation, the natural basis for the representation $\wedge^k = \wedge^k(\mathbb{C}^{mn})$ is parametrized by subsets of $[m] \times [n]$ with k elements.

Recall that, as a $U_q(gl_m) \otimes U_q(gl_n)$ -module, we have

$$\wedge^k(\mathbb{C}^{mn}) = \sum_{\lambda} V_{\lambda}(\mathbb{C}^m) \otimes V_{\lambda'}(\mathbb{C}^n)$$

where $|\lambda| = k$. Further, λ has at most m parts and λ' has at most n parts, that is, the shape λ fits inside the $m \times n$ ‘rectangle’.

For a shape $\lambda = (\lambda_1, \dots, \lambda_m)$ with $\lambda' = (\lambda'_1, \dots, \lambda'_n)$, consider the subset $c_{\lambda} \subset [mn]$ defined as:

$$c_{\lambda} = \left\{ \begin{array}{l} 1, m+1, \dots, m * (\lambda_1 - 1) + 1, \\ 2, m+2, \dots, m * (\lambda_2 - 1) + 2, \\ \dots \\ m, 2m, \dots, m * (\lambda_m - 1) \end{array} \right\}$$

Equivalently,

$$c_{\lambda} = \left\{ \begin{array}{l} 1, 2, \dots, \lambda'_1, \\ m+1, m+2, \dots, m + \lambda'_2, \\ \dots \\ m * (n-1) + 1, m * (n-1) + 2, \dots, m * (n-1) + \lambda'_n \end{array} \right\}$$

Under the identification of $[mn]$ with $[m] \times [n]$, we have

$$c_{\lambda} = \{(i, j) \mid 1 \leq i \leq \lambda'_j, 1 \leq j \leq \lambda_i\}$$

We slightly abuse the notation and write $(i, j) \in \lambda$ as a short-form for $(i, j) \in c_{\lambda}$.

With this notation, we have the following important lemma:

Lemma 34 *Consider the $U_q(gl_m) \otimes U_q(gl_n)$ -module $\wedge^k(\mathbb{C}^{mn})$. For a shape λ which fits in the $m \times n$ rectangle with $|\lambda| = k$, the weight vector $v_{c_{\lambda}} \in \wedge^k$ is the highest $U_q(gl_m) \otimes U_q(gl_n)$ -weight vector of weight (λ, λ') .*

Proof: The lemma follows from the observation that $E_i^L(v_{c_{\lambda}}) = E_j^R(v_{c_{\lambda}}) = 0$ for all i, j . Q.E.D.

Now we turn our attention to the construction of the $U_q(gl_m) \otimes U_q(gl_n)$ -equivariant map

$$\psi_a : \wedge^{a+1} \rightarrow \wedge^a \otimes \wedge^1$$

As a $U_q(gl_m) \otimes U_q(gl_n)$ -module, we have the following decomposition

$$\wedge^{a+1} = \sum_{\lambda: |\lambda|=a+1} V_{\lambda}(\mathbb{C}^m) \otimes V_{\lambda'}(\mathbb{C}^n)$$

Moreover, $v_{c_{\lambda}}$ is the highest-weight vector for the $U_q(gl_m) \otimes U_q(gl_n)$ -submodule $V_{\lambda}(\mathbb{C}^m) \otimes V_{\lambda'}(\mathbb{C}^n)$ of $\wedge^{a+1}$.

Thus, in order to construct the $U_q(gl_m) \otimes U_q(gl_n)$ -equivariant map ψ_a , we need to simply define the images $\psi_a(v_{c_\lambda})$ inside $\Lambda^a \otimes \Lambda^1$. Moreover the vector $\psi_a(v_{c_\lambda})$ should be a highest-weight vector of weight (λ, λ') . Note that, unlike Λ^{a+1} , $\Lambda^a \otimes \Lambda^1$ is not multiplicity-free. Below, we outline the construction of a highest-weight vector (upto scalar multiple) v_λ of weight (λ, λ') inside $\Lambda^a \otimes \Lambda^1$.

We begin with some notation. As before, fix a shape λ which fits in the $m \times n$ rectangle with $|\lambda| = a + 1$. Write $\lambda = (\lambda_1, \dots, \lambda_m)$ with $\lambda' = (\lambda'_1, \dots, \lambda'_n)$ and

$$c_\lambda = \{(i, j) \mid 1 \leq i \leq \lambda'_j, 1 \leq j \leq \lambda_i\}$$

For $(i, j) \in \lambda$, we set

$$\begin{aligned} t_{i,j} &= v_{c_\lambda - \{(i,j)\}} \in \Lambda^a \\ \chi_{i,j} &= v_{\{(i,j)\}} \in \Lambda^1 \end{aligned}$$

In other words, $t_{i,j}$ is the vector in Λ^a corresponding to the subset obtained from the subset c_λ by removing the element $(i, j) \in \lambda$. Further, $\chi_{i,j}$ is the vector in Λ^1 corresponding to the singleton set containing the element (i, j) . Below, we abuse notations and denote by $t_{i,j}$ and $\chi_{i,j}$ also the subsets that correspond to these vectors.

Lemma 35 *For $(i, j) \in \lambda$, $1 \leq k < m$, $1 \leq l < n$,*

- $E_k^L(t_{i,j}) = 0$ if $i \neq k$.
- $E_i^L(t_{i,j}) = t_{i+1,j}$ if $(i+1, j) \in \lambda$ and 0 otherwise.
- If $(i+1, j) \in \lambda$, $q^{-h_i^L}(t_{i+1,j}) = q^{\lambda_{i+1} - \lambda_i - 1} t_{i+1,j}$.
- $E_l^R(t_{i,j}) = 0$ if $j \neq l$.
- $E_j^R(t_{i,j}) = (-1)^{\lambda'_j - 1} q^{\lambda'_{j+1} - \lambda'_j} t_{i,j+1}$ if $(i, j+1) \in \lambda$ and 0 otherwise.
- If $(i, j+1) \in \lambda$, $q^{-h_j^R}(t_{i,j+1}) = q^{\lambda'_{j+1} - \lambda'_j - 1} t_{i,j+1}$.

Proof: Let $k \neq i$ and consider $E_k^L(t_{i,j})$. Note that, for all j' , if $(k+1, j') \in t_{i,j}$, then $(k, j') \in t_{i,j}$. Thus, by definition of E_k^L , we have $E_k^L(t_{i,j}) = 0$.

Now consider $E_i^L(t_{i,j})$. Note that $(i, j) \notin t_{i,j}$. If $(i+1, j) \in \lambda$, then $(i+1, j) \in t_{i,j}$. Further, for all $j' < j$, if $(i+1, j') \in t_{i,j}$ then $(i, j') \in t_{i,j}$. Thus, by definition $E_i^L(t_{i,j})$ operates only at the position $(i+1, j)$ if $(i+1, j) \in \lambda$ and produces the subset $t_{i+1,j}$.

Now we assume that $(i+1, j) \in \lambda$, and evaluate $q^{-h_i^L}(t_{i+1,j})$. Note that, except for $(i+1, j)$, $(i+1, j') \in t_{i+1,j}$ for $1 \leq j' \leq \lambda_{i+1}$. Also, for $j' > \lambda_{i+1}$, $(i+1, j') \notin t_{i+1,j}$. Thus $q^{\epsilon_{i+1}^L}(t_{i+1,j}) = q^{\lambda_{i+1} - 1}$. Similarly, $q^{\epsilon_i^L}(t_{i+1,j}) = q^{\lambda_i}$. Therefore,

$$q^{-h_i^L}(t_{i+1,j}) = q^{\lambda_{i+1} - \lambda_i - 1} t_{i+1,j}$$

It is easy to see that $E_l^R(t_{i,j}) = 0$ if $j \neq l$. So, we turn our attention to $E_j^R(t_{i,j})$. Note that, for i' such that $\lambda'_{j+1} < i' \leq \lambda'_j$, $(i', j) \in t_{i,j}$ and $(i', j+1) \notin t_{i,j}$. For other values of i' except i , either both or none of (i', j) and $(i', j+1)$ belong to $t_{i,j}$. Therefore, as expected, $E_j^R(t_{i,j})$ operates only at the position $(i, j+1)$ if $(i, j+1) \in \lambda$. Further, by definition of E_j^R , if $(i, j+1) \in \lambda$, we have

$$E_j^R(t_{i,j}) = (-1)^{\lambda'_j - 1} q^{\lambda'_{j+1} - \lambda'_j} t_{i,j+1} \text{ if } (i, j+1) \in t_{i,j}$$

The sign $(-1)^{\lambda'_j - 1}$ results from the fact that exactly $\lambda'_j - 1$ elements of $[mn]$ strictly in the range from (i, j) to $(i, j+1)$ belong to $t_{i,j}$.

We skip the proof for the last assertion as it follows from a similar reasoning applied earlier for the left E -operator. Q.E.D.

Lemma 36 For $(i, j) \in \lambda$,

•

$$E_i^L(t_{i,j} \otimes \chi_{i,j}) = \begin{cases} t_{i+1,j} \otimes \chi_{i,j} & \text{if } (i+1, j) \in \lambda \\ 0 & \text{otherwise} \end{cases}$$

• If $(i+1, j) \in \lambda$, then

$$E_i^L(t_{i+1,j} \otimes \chi_{i+1,j}) = q^{\lambda_{i+1}-\lambda_i-1} t_{i+1,j} \otimes \chi_{i,j}$$

•

$$E_j^R(t_{i,j} \otimes \chi_{i,j}) = \begin{cases} (-1)^{\lambda'_j-1} q^{\lambda'_{j+1}-\lambda'_j} t_{i,j+1} \otimes \chi_{i,j} & \text{if } (i, j+1) \in \lambda \\ 0 & \text{otherwise} \end{cases}$$

• If $(i, j+1) \in \lambda$, then

$$E_j^R(t_{i,j+1} \otimes \chi_{i,j+1}) = q^{\lambda'_{j+1}-\lambda'_j-1} t_{i,j+1} \otimes \chi_{i,j}$$

• For remaining $1 \leq k < m$ and $1 \leq l < n$, $E_k^L(t_{i,j} \otimes \chi_{i,j}) = E_l^R(t_{i,j} \otimes \chi_{i,j}) = 0$.

Proof: For the first assertion, consider

$$E_i^L(t_{i,j} \otimes \chi_{i,j}) = E_i^L(t_{i,j}) \otimes \chi_{i,j} + q^{-h_i^L} (t_{i,j}) \otimes E_i^L(\chi_{i,j})$$

As $(i+1, j) \notin \chi_{i,j}$, $E_i^L(\chi_{i,j}) = 0$. Therefore, the claim follows from the previous lemma.

For the second assertion, let us assume that $(i+1, j) \in \lambda$. Then

$$E_i^L(t_{i+1,j} \otimes \chi_{i+1,j}) = E_i^L(t_{i+1,j}) \otimes \chi_{i+1,j} + q^{-h_i^L} (t_{i+1,j}) \otimes E_i^L(\chi_{i+1,j})$$

Note that, from the previous lemma $E_i^L(t_{i+1,j}) = 0$. Also, $E_i^L(\chi_{i+1,j}) = \chi_{i,j}$. Again, using the previous lemma, we have

$$E_i^L(t_{i+1,j} \otimes \chi_{i+1,j}) = q^{\lambda_{i+1}-\lambda_i-1} t_{i+1,j} \otimes \chi_{i,j}$$

The third and fourth assertions are proved in a similar fashion. Q.E.D.

Lemma 37 Let $v_\lambda \in \wedge^a \otimes \wedge^1$ be defined as follows:

$$v_\lambda = \sum_{(k,l) \in \lambda} \alpha_{k,l} t_{k,l} \otimes \chi_{k,l}$$

where

$$\alpha_{k,l} = (-1)^{\lambda'_1 + \dots + \lambda'_{l-1} + k} q^{k+l-\lambda_k}$$

Then v_λ is a highest-weight vector of weight (λ, λ') .

Proof: It is clear that v_λ is a weight vector of weight (λ, λ') . Below, we show that it is a highest-weight vector by checking that $E_i^L(v_\lambda) = E_j^R(v_\lambda) = 0$ for all i, j .

Towards this, by previous lemma, we have

$$\begin{aligned} E_i^L(v_\lambda) &= \sum_{(k,l) \in \lambda} \alpha_{k,l} E_i^L(t_{k,l} \otimes \chi_{k,l}) \\ &= \sum_{(i,l) \in \lambda} \alpha_{i,l} E_i^L(t_{i,l} \otimes \chi_{i,l}) + \sum_{(i+1,l) \in \lambda} \alpha_{i+1,l} E_i^L(t_{i+1,l} \otimes \chi_{i+1,l}) \\ &= \sum_{l: (i,l) \& (i+1,l) \in \lambda} (\alpha_{i,l} E_i^L(t_{i,l} \otimes \chi_{i,l}) + \alpha_{i+1,l} E_i^L(t_{i+1,l} \otimes \chi_{i+1,l})) \end{aligned}$$

For l such that both (i, l) and $(i+1, l)$ are in λ , from previous lemma, we have

$$\begin{aligned} E_i^L(t_{i,l} \otimes \chi_{i,l}) &= t_{i+1,l} \otimes \chi_{i,l} \\ E_i^L(t_{i+1,l} \otimes \chi_{i+1,l}) &= q^{\lambda_{i+1}-\lambda_i-1} t_{i+1,l} \otimes \chi_{i,l} \end{aligned}$$

Therefore, the coefficient of $t_{i+1,l} \otimes \chi_{i,l}$ in $E_i^L(v_\lambda)$ is

$$\begin{aligned}
&= \alpha_{i,l} + q^{\lambda_{i+1}-\lambda_i-1} \alpha_{i+1,l} \\
&= (-1)^{\lambda'_1+\dots+\lambda'_{i-1}+i} q^{i+l-\lambda_i} + q^{\lambda_{i+1}-\lambda_i-1} (-1)^{\lambda'_1+\dots+\lambda'_{i-1}+i+1} q^{i+1+l-\lambda_{i+1}} \\
&= (-1)^{\lambda'_1+\dots+\lambda'_{i-1}+i} (q^{i+l-\lambda_i} - q^{i+l-\lambda_{i+1}}) \\
&= 0
\end{aligned}$$

Thus, $E_i^L(v_\lambda) = 0$. A similar analysis shows that, the coefficient of $t_{k,j+1} \otimes \chi_{k,j}$ in $E_j^R(v_\lambda)$ is

$$\begin{aligned}
&= \alpha_{k,j} (-1)^{\lambda'_j-1} q^{\lambda'_{j+1}-\lambda'_j} + \alpha_{k,j+1} q^{\lambda'_{j+1}-\lambda'_j-1} \\
&= (-1)^{\lambda'_1+\dots+\lambda'_{j-1}+k} q^{k+j-\lambda_k} (-1)^{\lambda'_j-1} q^{\lambda'_{j+1}-\lambda'_j} + (-1)^{\lambda'_1+\dots+\lambda'_j+k} q^{k+j+1-\lambda_k} q^{\lambda'_{j+1}-\lambda'_j-1} \\
&= 0
\end{aligned}$$

This shows that $E_j^R(v_\lambda) = 0$ and hence establishes the claim that v_λ is a highest-weight vector in $\wedge^a \otimes \wedge^1$. Q.E.D.

We remark that in the above expression for v_λ , the coefficient, $\alpha_{1,1}$, of the term $t_{1,1} \otimes \chi_{1,1}$ has the least q -degree. We may normalize v_λ so as to ensure that $\alpha_{1,1} = 1$ and all the other terms have strictly positive q -degree.

Now we are ready to define the $U_q(gl_m) \otimes U_q(gl_n)$ -equivariant map

$$\psi_a : \wedge^{a+1} \rightarrow \wedge^a \otimes \wedge^1$$

This is done by simply setting $\psi_a(v_{e_\lambda}) = v_\lambda$. It is easily seen that there is a unique $U_q(gl_m) \otimes U_q(gl_n)$ -equivariant extension of ψ_a to all of $\wedge^{a+1}$. Moreover, this extension matches the classical $U_1(gl_{mn})$ -equivariant construction at $q = 1$.

Next, we prepare towards the construction of the $U_q(gl_m) \otimes U_q(gl_n)$ -equivariant map ψ'_a .

Lemma 38 For $(i, j) \in \lambda$,

- $$E_i^L(\chi_{i,j} \otimes t_{i,j}) = \begin{cases} q^{-1} \chi_{i,j} \otimes t_{i+1,j} & \text{if } (i+1, j) \in \lambda \\ 0 & \text{otherwise} \end{cases}$$
- If $(i+1, j) \in \lambda$, then $E_i^L(\chi_{i+1,j} \otimes t_{i+1,j}) = \chi_{i,j} \otimes t_{i+1,j}$
- $$E_j^R(\chi_{i,j} \otimes t_{i,j}) = \begin{cases} (-1)^{\lambda'_j-1} q^{\lambda'_{j+1}-\lambda'_j-1} \chi_{i,j} \otimes t_{i,j+1} & \text{if } (i, j+1) \in \lambda \\ 0 & \text{otherwise} \end{cases}$$
- If $(i, j+1) \in \lambda$, then
$$E_j^R(\chi_{i,j+1} \otimes t_{i,j+1}) = \chi_{i,j} \otimes t_{i,j+1}$$
- For remaining $1 \leq k < m$ and $1 \leq l < n$, $E_k^L(\chi_{i,j} \otimes t_{i,j}) = E_l^R(\chi_{i,j} \otimes t_{i,j}) = 0$.

Proof: For the first assertion, consider

$$E_i^L(\chi_{i,j} \otimes t_{i,j}) = E_i^L(\chi_{i,j}) \otimes t_{i,j} + q^{-h_i^L}(\chi_{i,j}) \otimes E_i^L(t_{i,j})$$

As $(i+1, j) \notin \lambda$, $E_i^L(\chi_{i,j}) = 0$. Further, $q^{-h_i^L}(\chi_{i,j}) = q^{-1} \chi_{i,j}$. Therefore, the claim follows.

For the second assertion, let us assume that $(i+1, j) \in \lambda$. Then

$$E_i^L(\chi_{i+1,j} \otimes t_{i+1,j}) = E_i^L(\chi_{i+1,j}) \otimes t_{i+1,j} + q^{-h_i^L}(\chi_{i+1,j}) \otimes E_i^L(t_{i+1,j})$$

Note that, $E_i^L(t_{i+1,j}) = 0$. Also, $E_i^L(\chi_{i+1,j}) = \chi_{i,j}$. Therefore, we have

$$E_i^L(\chi_{i+1,j} \otimes t_{i+1,j}) = \chi_{i,j} \otimes t_{i+1,j}$$

For the third assertion, consider

$$\begin{aligned} E_j^R(\chi_{i,j} \otimes t_{i,j}) &= E_j^R(\chi_{i,j}) \otimes t_{i,j} + q^{-h_j^R}(\chi_{i,j}) \otimes E_j^R(t_{i,j}) \\ &= q^{-1}\chi_{i,j} \otimes E_j^R(t_{i,j}) \end{aligned}$$

Recall that, we have

$$E_j^R(t_{i,j}) = (-1)^{\lambda'_j-1} q^{\lambda'_{j+1}-\lambda'_j} t_{i,j+1} \text{ if } (i, j+1) \in \lambda \text{ and } 0 \text{ otherwise}$$

Therefore, the claim follows.

For the fourth claim, we assume $(i, j+1) \in \lambda$. Then

$$\begin{aligned} E_j^R(\chi_{i,j+1} \otimes t_{i,j+1}) &= E_j^R(\chi_{i,j+1}) \otimes t_{i,j+1} + q^{-h_j^R}(\chi_{i,j+1}) \otimes E_j^R(t_{i,j+1}) \\ &= \chi_{i,j} \otimes t_{i,j+1} \end{aligned}$$

The last claim can be easily proved. Q.E.D.

Lemma 39 Let $v_\lambda \in \wedge^1 \otimes \wedge^a$ be defined as follows:

$$v_\lambda = \sum_{(k,l) \in \lambda} \beta_{k,l} \chi_{k,l} \otimes t_{k,l}$$

where

$$\beta_{k,l} = (-1)^{\lambda'_1 + \dots + \lambda'_{i-1} + k} q^{\lambda'_i - k - l}$$

Then v_λ is a highest-weight vector of weight (λ, λ') .

Proof: Clearly, v_λ is a weight-vector of weight (λ, λ') . We now check that $E_i^L(v_\lambda) = 0$ for all i . As expected, this finally reduces to checking if the following expression, coefficient of $\chi_{i,l} \otimes t_{i+1,l}$ in $E_i^L(v_\lambda)$, is zero. Towards this, consider

$$\begin{aligned} &= q^{-1}\beta_{i,l} + \beta_{i+1,l} \\ &= q^{-1}(-1)^{\lambda'_1 + \dots + \lambda'_{i-1} + i} q^{\lambda'_i - i - l} + (-1)^{\lambda'_1 + \dots + \lambda'_{i-1} + i+1} q^{\lambda'_i - i - 1 - l} \\ &= 0 \end{aligned}$$

Similarly, to check if $E_j^R(v_\lambda) = 0$, we need to check if the following expression, coefficient of $\chi_{k,j} \otimes t_{k,j+1}$ in $E_j^R(v_\lambda)$, is zero. Towards this, consider

$$\begin{aligned} &= \beta_{k,j}(-1)^{\lambda'_j-1} q^{\lambda'_{j+1}-\lambda'_j-1} + \beta_{k,j+1} \\ &= (-1)^{\lambda'_1 + \dots + \lambda'_{j-1} + k} q^{\lambda'_j - k - j} (-1)^{\lambda'_j-1} q^{\lambda'_{j+1}-\lambda'_j-1} + (-1)^{\lambda'_1 + \dots + \lambda'_j + k} q^{\lambda'_{j+1} - k - j - 1} \\ &= 0 \end{aligned}$$

Thus, we have verified that $E_i^L(v_\lambda) = E_j^R(v_\lambda) = 0$ for all i, j . This shows that v_λ is a highest-weight vector. Q.E.D.

Now we are ready to define the $U_q(gl_m) \otimes U_q(gl_n)$ -equivariant map

$$\psi'_a : \wedge^{a+1} \rightarrow \wedge^1 \otimes \wedge^a$$

As expected, this is done by simply setting $\psi'_a(v_{c_\lambda}) = v'_\lambda$ and taking the unique $U_q(gl_m) \otimes U_q(gl_n)$ -equivariant extension. Also, as before, this extension matches the classical $U_1(gl_{mn})$ -equivariant construction at $q = 1$.

Note that $\wedge^{a+1}$ and $\wedge^1 \otimes \wedge^a$ have normal bases. Whence, by Prop. 33, there is the $U_q(gl_m) \otimes U_q(gl_n)$ -equivariant map:

$$\psi'^*_a : \wedge^1 \otimes \wedge^a \rightarrow \wedge^{a+1}$$

Finally, we construct $\psi_{a,b}$ as follows:

$$\psi_{a,b} : \wedge^{a+1} \otimes \wedge^{b-1} \xrightarrow{\psi_a \otimes I_{\wedge^{b-1}}} \wedge^a \otimes \wedge^1 \otimes \wedge^{b-1} \xrightarrow{I_{\wedge^a} \otimes \psi'^*_{b-1}} \wedge^a \otimes \wedge^b$$

5 The crystal basis for $\wedge^K$

In this section we examine the crystal structure (see [9, 10]) of the $U_q(gl_m) \otimes U_q(gl_n)$ -module $\wedge^K(\mathbb{C}^{m \times n})$. We show that there is a sign function $sign^*$ on K -subsets of $[mn]$ such that the collection $\mathcal{B}^* = \{sign^*(c) \cdot v_c\}_c$ is a crystal basis for $\wedge^K$.

We identify $[mn]$ with $[m] \times [n]$ and also order the elements as follows:

$$(1, 1) \prec (2, 1) \prec \dots (m, 1) \prec (1, 2) \prec \dots (m-1, n) \prec (m, n)$$

In other words $(i, j) \prec (i', j')$ iff either $j < j'$ or $j = j'$ with $i < i'$. For $(i, j) \prec (i', j')$, we denote by $[(i, j), (i', j')]$ as the indices between (i, j) and (i', j') including both (i, j) and (i', j') .

Recall that (cf. Section 2), as a $\mathbb{C}(q)$ -vector space, $\wedge^K(\mathbb{C}^{mn})$ is generated by the basis vectors $\mathcal{B} = \{v_c | c \subseteq [mn], |c| = K\}$. Let us fix an index i and look at the sub-algebra U_i^L of $U_q(gl_m)$ generated by E_i^L, F_i^L and h_i^L . We define the standard $U_q(sl_2)$ generated by symbols e, f, h satisfying the following equations:

$$\begin{aligned} q^h q^{-h} &= 1 \\ q^h e q^{-h} &= q^2 e \\ q^h f q^{-h} &= q^{-2} f \\ ef - fe &= \frac{e^h - e^{-h}}{q - q^{-1}} \end{aligned}$$

We use the Hopf Δ :

$$\Delta q^h = q^h \otimes q^h, \Delta e = e \otimes 1 + q^{-h} \otimes e, \Delta f = f \otimes q^h + 1 \otimes f$$

In other words, they satisfy exactly the same relations that e_i^L, f_i^L, h_i^L satisfy, including the Hopf. Clearly, U_i^L is isomorphic to $U_q(sl_2)$ as algebras and we denote this isomorphism by $L : U_i^L \rightarrow U_q(sl_2)$.

We construct the $U_q(sl_2)$ -module $\mathbb{C}^2$ with basis x_1, x_2 with the action:

$$ex_2 = x_1, ex_1 = 0, fx_2 = 0, fx_1 = x_2, q^h x_1 = qx_1, q^h x_2 = q^{-1}x_2$$

With the Hopf Δ above, $M = \otimes_{i=1}^N \mathbb{C}^2$ is a $U_q(sl_2)$ -module with the basis $\mathcal{S} = \{y_1 \otimes \dots \otimes y_N | y_i \in \{x_1, x_2\}\}$, and with the action:

$$e(y_1 \otimes \dots \otimes y_N) = \sum_j \left(\prod_{k=1}^{j-1} q^{-h}(y_k) \right) \cdot y_1 \otimes \dots \otimes y_{j-1} \otimes e(y_j) \otimes y_{j+1} \otimes \dots \otimes y_N$$

A similar expression may be written for the action of f .

Let us identify $[mn]$ with $[m] \times [n]$ and define the **signature** $\sigma_i^L(c)$, for $c \subseteq [mn]$. Towards this, we define

$$\begin{aligned} I(c) &= \{1 \leq j \leq n \mid \text{both } (i, j), (i+1, j) \in c\} \\ J(c) &= \{1 \leq j \leq n \mid \text{both } (i, j), (i+1, j) \notin c\} \\ S(c) &= \{(i', j') \in c \mid i' \neq i \text{ and } i' \neq i+1\} \end{aligned}$$

The signature $\sigma_i^L(c)$ is the tuple $(I(c), J(c), S(c))$.

Next, for a $\sigma = (I, J, S)$, we define the vector space $V_{\sigma, i}^L$ as the $\mathbb{C}(q)$ -span of all elements

$$\mathcal{B}_{\sigma, i}^L = \{v_c \mid \sigma_i^L(c) = \sigma\}$$

Let $N = n - |I| - |J|$ and let $M = \otimes^N \mathbb{C}^2$ be the $U_q(sl_2)$ -module as above.

We prove the following:

Proposition 40 *Given $\sigma = (I_\sigma, J_\sigma, S_\sigma)$ as above,*

(i) $V_{\sigma, i}^L$ is a U_i^L -invariant subspace.

(ii) The $U_q(sl_2)$ module M is isomorphic to the U_i^L -module $V_{\sigma,i}^L$ via the isomorphism L above.

Proof: For any $v_c \in \mathcal{B}_{\sigma,i}^L$, if $E_i^L(v_c) = \sum \alpha(c') \cdot v_{c'}$, then it is clear that $v_{c'} \in \mathcal{B}_{\sigma,i}^L$ as well. The same holds for F_i^L and h_i^L . This proves (i) above. For (ii), first note that

$$E_i^L = \sum_j \left(\prod_{k=1}^{j-1} q^{-h_{(k-1)m+i}} \right) e_{(j-1)m+i}$$

which matches the Hopf Δ of $U_q(sl_2)$. Next, if $j \in I(c) \cup J(c)$ then the index j is irrelevant to the action of E_i^L on v_c , whence in the restriction to $V_{\sigma,i}^L$, the indices in $I_\sigma \cup J_\sigma$ do not play a role.

Next, note that $|\mathcal{B}_{\sigma,i}^L| = 2^N$. Assume for simplicity that $I_\sigma \cup J_\sigma = \{N+1, \dots, n\}$. Indeed, we may set up a $U_q(sl_2)$ -module isomorphism ι_L by setting

$$\iota_L(v_c) = y_1 \otimes \dots \otimes y_N \text{ such that } y_k = \begin{cases} x_1 & \text{iff } (i, k) \in c \\ x_2 & \text{otherwise} \end{cases}$$

One may verify that $\iota_L : V_{\sigma,i}^L \rightarrow M$ is indeed equivariant via L . Q.E.D.

Proposition 41 *The elements $\mathcal{B}$ is a crystal basis for $\wedge^K(\mathbb{C}^{mn})$ for the action of $U_q(gl_m)$.*

Proof: This is obtained by first noting that $\mathcal{S}$ is indeed a crystal basis for M , see [9], for example. Next, the equivariance of ι_L shows that for $v_c \in \mathcal{B}_{\sigma,i}^L$,

$$\tilde{E}_i^L(v_c) = \iota_L^{-1}(\tilde{e}(\iota_L(v_c)))$$

This proves that $\mathcal{B}_{\sigma,i}^L$ is indeed a crystal basis for $V_{\sigma,i}^L$. Next, by applying Proposition 40 for all i and all σ , we see that $\{\mathcal{B}_{\sigma,i}^L | i, \sigma\}$ together cover $\mathcal{B}$. Q.E.D.

We now move to the trickier $U_q(gl_n)$ -action. Let us denote by $\epsilon_{i,j}$ the weight $\epsilon_{(j-1)*m+i}$ and $h_{i,j} = \epsilon_{i,j} - \epsilon_{i,j+1}$. There are two sources of complications.

- The operator E_k^R may be re-written as:

$$E_k^R = \sum_i \prod_{a=i+1}^m (q^{-h_{a,k}}) E_{(k-1)m+i, km+i-1} = \sum_i E_{(k-1)m+i, km+i-1} \left(\prod_{a=i+1}^m q^{-h_{a,k}} \right)$$

Thus, the Hopf works from the “right”.

- For a general v_c , if $E_{(k-1)m+i, km+i-1} v_c$ is non-zero then it is $\pm v_d$, where $v_d = v_c - (i, k+1) + (i, k)$ where the sign is $(-1)^M$ where M is the number of elements in $c \cap [(i+1, k), \dots, (i-1, k+1)]$.

To fix the sign, we first define an “intermediate global” sign as follows. For a set $c \subset [m] \times [n]$, we define $c^* \subset [m] \times [n]$ as that obtained by moving the elements of c to the right, as far as they can go (see Example 46). Note that $F_k^R(c^*) = 0$ for all k and thus c^* is *one of the* lowest weight vectors in $\wedge^K(\mathbb{C}^{mn})$. For an $(i, j) \in c$, let (i, j^*) be its final position in c^* . We may define j^* explicitly as $n - |\{j' | (i, j') \in c, j' > j\}|$. Next, we define for $(i, j) \in c$,

$$\begin{aligned} S_{i,j}(c) &= \{(i', j') \in c \mid (i, j) \prec (i', j') \prec (i', j^*) \prec (i, j^*)\} \\ n_{ij} &= |S_{i,j}(c)| \end{aligned}$$

Setting $N_c = \sum_{(i,j) \in c} n_{ij}$ we finally define:

$$\begin{aligned} \text{sign}(c) &= (-1)^{N_c} \\ \text{sign}(d/c) &= \text{sign}(d) / \text{sign}(c) \end{aligned}$$

Lemma 42 Let $v_c \in \mathcal{B}_{\sigma,k}^R$ be such that $E_{(k-1)m+i, km+i-1} v_c \neq 0$, then

$$E_{(k-1)m+i, km+i-1} v_c = \text{sign}(d/c) v_d$$

where $v_d = v_c - (i, k+1) + (i, k)$.

Proof: It is clear that $c^* = d^*$ and thus for $(i, k+1) \in c$ and $(i, k) \in d$, let (i, k^*) be the final position of both $(i, k+1) \in c$ and $(i, k) \in d$. For $(i, k+1) \prec (i', j')$ or $(i', j') \prec (i, k)$ we have (i) $S_{i', j'}(c) = S_{i', j'}(d)$ and (ii) $(i', j') \in S_{i, k+1}(c)$ iff $(i', j') \in S_{i, k}(d)$.

Next, it is clear that (i) $S_{i, k}(d) \supseteq S_{i, k+1}(c)$, and (ii) for $(i, k) \prec (i', j') \prec (i, k+1)$, $S_{i', j'}(d) \subseteq S_{i', j'}(c)$ and in fact, $S_{i', j'}(c) - S_{i', j'}(d)$ can at most be the element $(i, k+1)$.

Now let us look at $S_{i, k}(d) - S_{i, k+1}(c)$. These contain all $(i', j') \in c$ such that

$$(i, k) \prec (i', j') \prec (i, k+1) \prec (i', j'^*) \prec (i, k^*)$$

On the other hand, for $(i', j') \in c$ such that $(i, k) \prec (i', j') \prec (i, k+1)$, which are not counted above, it must be that $(i, k^*) \prec (i', j'^*)$ in which case, $S_{i', j'}(c) = S_{i', j'}(d) \cup \{(i, k+1)\}$.

In short, for every $(i', j') \in c$ such that $(i, k) \prec (i', j') \prec (i, k+1)$ either it contributes to an increment in $S_{i, k}(d)$ over $S_{i, k+1}(c)$ or a decrement in $S_{i', j'}(d)$ over $S_{i', j'}(c)$. Ofcourse, the two cases are exclusive.

Thus we have $\text{sign}(d)/\text{sign}(c) = (-1)^M$ where M is exactly the number of elements in $c \cap [(i+1, k), \dots, (i-1, k+1)]$. Q.E.D.

Next, we define a new Hopf Δ' on $U_q(sl_2)$ as

$$\Delta' q^h = q^h \otimes q^h, \Delta' e = 1 \otimes e + e \otimes q^{-h}, \Delta' f = q^h \otimes f + f \otimes 1$$

We denote by M' , the $U_q(sl_2)$ -module $\otimes^N \mathbb{C}^2$ via the Hopf Δ' and with the basis $\mathcal{S} = \{y_1 \otimes \dots \otimes y_N | y_i \in \{x_1, x_2\}\}$. Under Δ' we have:

$$e(y_1 \otimes \dots \otimes y_N) = \sum_j \left(\prod_{k=j+1}^N q^{-h}(y_k) \right) \cdot y_1 \otimes \dots \otimes y_{j-1} \otimes e(y_j) \otimes y_{j+1} \otimes \dots \otimes y_N$$

We denote by U_k^R the algebra generated by E_k^R, F_k^R, h_k^R and let $R : U_k^R \rightarrow U_q(sl_2)$ be the natural isomorphism.

As before, we define $\sigma_k^R(c)$ analogously as

$$\begin{aligned} I(c) &= \{1 \leq i \leq m \mid \text{both } (i, k), (i, k+1) \in c\} \\ J(c) &= \{1 \leq i \leq m \mid \text{both } (i, k), (i, k+1) \notin c\} \\ S(c) &= \{(i', k') \in c \mid k' \neq k \text{ and } k' \neq k+1\} \end{aligned}$$

Next, for a $\sigma = (I, J, S)$, we define the vector space $V_{\sigma, k}^R$ as the $\mathbb{C}(q)$ -span of all elements

$$\mathcal{B}_{\sigma, k}^R = \{v_c \mid \sigma_k^R(c) = \sigma\}$$

Again, as before, let $N = n - |I| - |J|$. Let us also assume, for simplicity that $I \cup J = \{N+1, \dots, m\}$.

Proposition 43 Given σ as above,

(i) $V_{\sigma, k}^R$ is a U_k^R -invariant subspace.

(ii) The $U_q(sl_2)$ module M' is isomorphic to the U_k^R -module $V_{\sigma, k}^R$ via the isomorphism R above.

Proof: Part (i) above is obvious. For (ii), note that

$$E_k^R = \sum_i E_{(k-1)m+i, km+i-1} \left(\prod_{a=i+1}^m q^{-h_{a,k}} \right)$$

which matches the Hopf Δ' of $U_q(sl_2)$. Again, if $j \in I(c) \cup J(c)$ then the index j is irrelevant to the action of E_k^R on v_c , whence in the restriction to $V_{\sigma,k}^R$, the indices in $I \cup J$ do not play a role.

Next, note that $|\mathcal{B}_{\sigma,k}^R| = 2^N$. Recall that, we have assumed that $I \cup J = \{N+1, \dots, m\}$. Indeed, we may set up a $U_q(sl_2)$ -module isomorphism ι_R by setting

$$\iota_R(v_c) = \text{sign}(c) \cdot y_1 \otimes \dots \otimes y_N \text{ such that } y_i = \begin{cases} x_1 & \text{iff } (i, k) \in c \\ x_2 & \text{otherwise} \end{cases}$$

One may verify (using Lemma 42) that $\iota_R : V_{\sigma,k}^R \rightarrow M'$ is indeed equivariant via R . Q.E.D.

Proposition 44 *Let $\mathcal{B}' = \{\text{sign}(b) \cdot v_b | b \in \mathcal{B}\}$ be “signed” elements. Then the elements $\mathcal{B}'$ is a crystal basis for $\wedge^K(\mathbb{C}^{mn})$ for the action of $U_q(gl_n)$. In other words $\tilde{E}_k^R(v_c) = \pm v_d \cup 0$.*

Proof: Let $\mathcal{B}'_{\sigma,k}^R$ be the “signed” elements of $\mathcal{B}_{\sigma,k}^R$. We first note that $\mathcal{S}$ continues to be a crystal basis for M' . Next, the equivariance of ι_R shows that for $v_c \in \mathcal{B}'_{\sigma,k}^R$,

$$\tilde{E}_k^R(v_c) = \iota_R^{-1}(\tilde{e}(\iota_R(v_c)))$$

This proves that $\mathcal{B}'_{\sigma,k}^R$ is indeed a crystal basis for $V_{\sigma,k}^R$.

Thus, keeping in mind that the signs are allotted by our global *sign*-function and, by considering all σ and all k , we obtain the assertion. Q.E.D.

We now define our final global sign $\text{sign}^*(b)$ as follows. Firstly, let $S = \{b \mid F_i^L v_b = F_k^R v_b = 0\}$. These are the lowest weight vectors for both the left and the right action. We see that:

- For any $b \in S$, we have $b^* = b$.
- If $wt_i(b)$ denotes the cardinality of the set $\{(i, k) | (i, k) \in b\}$, then $wt_1(b) \leq \dots \leq wt_m(b)$.

We define $\text{sign}^*(b) = \text{sign}(b)$ for all b such that $b^* \in S$. Next, for a c such that $c^* \notin S$, we inductively (by (wt_i) above) define $\text{sign}^*(c) = \text{sign}^*(\tilde{F}_i^L(v_c))$ where $\tilde{F}_i^L(v_c) \neq 0$. By the commutativity of $\tilde{F}_i^L$ with $\tilde{F}_k^R$, we see that $\text{sign}^*(c)$ is well defined over all K -subsets of $[m] \times [n]$.

Let $v_b^* = \text{sign}^*(b) \cdot v_b$ and let $\mathcal{B}^* = \{v_b^* | v_b \in \mathcal{B}\}$.

Proposition 45 *The elements $\mathcal{B}^*$ is a crystal basis for $\wedge^K(\mathbb{C}^{mn})$ for the action of both $U_q(gl_n)$ and $U_q(gl_m)$. In other words $\tilde{E}_k^R(v_c^*) \in \mathcal{B}^* \cup 0$. and $\tilde{E}_i^L(v_c^*) \in \mathcal{B}^* \cup 0$.*

Proof: The proof follows from the commutativity condition and the well-defined-ness of sign^* . Q.E.D.

Example 46 *Let us consider $\wedge^2(\mathbb{C}^{2 \times 2})$ whose six elements, their matrix notation, and signs are given below:*

c	$matrix$	c^*	$sign^*(c^*)$	$sign^*(c)$
$\begin{array}{ c } \hline 1 \\ \hline 2 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 1 & 0 \\ \hline 1 & 0 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 0 & 1 \\ \hline 0 & 1 \\ \hline \end{array}$	1	1
$\begin{array}{ c } \hline 1 \\ \hline 3 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 1 & 1 \\ \hline 0 & 0 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 1 & 1 \\ \hline 0 & 0 \\ \hline \end{array}$	1	1
$\begin{array}{ c } \hline 1 \\ \hline 4 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 1 & 0 \\ \hline 0 & 1 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 0 & 1 \\ \hline 0 & 1 \\ \hline \end{array}$	1	1
$\begin{array}{ c } \hline 2 \\ \hline 3 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 0 & 1 \\ \hline 1 & 0 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 0 & 1 \\ \hline 0 & 1 \\ \hline \end{array}$	1	-1
$\begin{array}{ c } \hline 2 \\ \hline 4 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 0 & 0 \\ \hline 1 & 1 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 0 & 0 \\ \hline 1 & 1 \\ \hline \end{array}$	1	1
$\begin{array}{ c } \hline 3 \\ \hline 4 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 0 & 1 \\ \hline 0 & 1 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 0 & 1 \\ \hline 0 & 1 \\ \hline \end{array}$	1	1

For a $b \subseteq [m] \times [n]$ define the **left word** $LW(b)$ as the i -indices of all elements $(i, k) \in b$, read bottom to top within a column, reading the columns left to right. Similarly, define the **right word** $RW(b)$ as the k -indices of all elements $(i, k) \in b$, read right to left within a row, reading the rows from bottom to top. For a word w , let $rs(w)$ be the Robinson-Schenstead tableau associated with w , when read from left to right. Define the **left tableau** $LT(b) = rs(LW(b))$ and the **right tableau** as $RT(b) = rs(RW(b))$.

Example 47 Let $m = 3$ and $n = 4$ and let $b = \{1, 3, 5, 6, 9, 10, 11\}$.

$$\begin{array}{|c|c|c|c|} \hline 1 & 0 & 0 & 1 \\ \hline 0 & 1 & 0 & 1 \\ \hline 1 & 1 & 1 & 0 \\ \hline \end{array}
\quad
LW(b) = 3132321
\quad
RW(b) = 3214241$$

$$LT(b) = \begin{array}{|c|c|c|} \hline 1 & 1 & 2 \\ \hline 2 & 3 & 3 \\ \hline 3 & & \\ \hline \end{array}
\quad
RT(b) = \begin{array}{|c|c|c|} \hline 1 & 1 & 4 \\ \hline 2 & 2 & \\ \hline 3 & 4 & \\ \hline \end{array}$$

For semi-standard tableau, recall the crystal operators $\tilde{e}_i^T, \tilde{f}_i^T$, see for example, [9]. These crystal operators may be connected to our crystal operators via the following proposition. This obtains the result in [2].

Proposition 48 For any $v_b^* \in \mathcal{B}^*$ the crystal basis for $\wedge^K(\mathbb{C}^{mn})$ as above, we have:

- If $\tilde{E}_i^L(v_b^*) = v_c^*$ then $\tilde{e}_i^T(LT(b)) = LT(c)$.
- If $\tilde{E}_k^R(v_b^*) = v_c^*$ then $\tilde{e}_k^T(RT(b)) = RT(c)$.

A similar assertion holds for the $\tilde{F}$ -operators.

References

- [1] M. Artin, W. Schelter, J. Tate: Quantum deformations of GL_n , Commun. Pure. Appl. Math. 44 (1991), 879-895.

- [2] V. I. Danilov, G. A. Koshevoi: Bi-crystals and crystal $(GL(V), GL(W))$ duality, Preprint RIMS-1485, Kyoto Univ., Kyoto (2004).
- [3] E. Date, M. Jimbo, T. Miwa, Representations of $U_q(gl(n, \mathbb{C}))$ at $q = 0$ and the Robinson-Schensted correspondence, in Physics and Mathematics of strings, World Scientific, Singapore (1990).
- [4] V. Drinfeld: Quantum groups, in proceedings of the International Congress of Mathematicians (A. M. Gleason, ed), Amer. Math. Soc., Providence, RI (1986), 254-258.
- [5] W. Fulton, J. Harris: Representation theory, Springer Verlag (1991).
- [6] M. Jimbo: a q -analogue of $U(\mathcal{G})$ and the Yang-Baxter equation, Lett. Math. Phys. 10 (1985), 63-69.
- [7] T. Hayashi: Nonexistence of homomorphisms between quantum groups, Tokyo J. Math. 15 (1992), 431-435.
- [8] B. Leclerc and J. Thibon: The Robinson-Schensted correspondence as the quantum straightening at $q=0$, Electronic Journal of Combinatorics, Vol. 3, no. 2, (1996).
- [9] M. Kashiwara: Crystallizing the q -analogue of universal enveloping algebra, Commun. Math. Phys. 133 (1990) 249-260.
- [10] M. Kashiwara: On crystal bases of the q -analogue of the universal enveloping algebras, Duke Math. J. 63 (1991), 465-516.
- [11] A. Klimyk, K. Schmüdgen, Quantum groups and their representations, Springer (1997).
- [12] G. Lusztig, Canonical bases in tensor products, Proc. Nat. Acad. Sci. USA, vo. 89 (1992), 8177-8179.
- [13] K. Mulmuley, M. Sohoni: Geometric complexity theory: An approach to the P vs. NP and related problems, SIAM J. comput. vol. 31, no. 2, (2001), 496-526.
- [14] K. Mulmuley, M. Sohoni: Geometric complexity theory II: towards explicit obstructions for embeddings among class varieties, arXiv cs.CC/0612134, December, 2006, also in SIAM J. Comput. 38(3): 1175-1206 (2008)
- [15] K. Mulmuley, M. Sohoni: Geometric Complexity Theory IV: quantum group for the Kronecker problem, arXiv:cs/0703110 (2007).
- [16] L. G. Valiant: Completeness classes in algebra, Proc. 11th ACM STOC (1979), 249-261.